

Product datasheet for **TP301323L**

Morg1 (WDR83) (NM_032332) Human Recombinant Protein


Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human mitogen-activated protein kinase organizer 1 (MORG1), transcript variant 2, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201323 protein sequence Red =Cloning site Green =Tags(s) MAFPEPKPRPPELPQKRLKTLDCGQGAVRAVRFNVDGNYCLTCGSDKTLKLWNPLRGTLRLRTYSGHGYEV LDAAGSFDNSSLCSGGGDKAVVLWDVASGQVVRKFRGHAGKVNTVQFNEEATVILSGSIDSSIRCWDCRS RRPEPVQTLDEARDGVSSVKVSDHEILAGSVDGRVRRYDLRMGQLFSDYVGPITCTCFSRDGGQCTLVSS LDSTLRLLDKDTGELLGEYKGHKNQEYKLDCCLSERDTHVSCSEDGKVFFWDLVEGALALALPVGSGVV QSLAYHPTPECLLTAMGGSVQCWREEAYEAEDGAG TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	34.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_115708</u>
Locus ID:	84292


[View online »](#)

UniProt ID:	<u>Q9BRX9</u>
RefSeq Size:	1219
Cytogenetics:	19p13.13
RefSeq ORF:	945
Synonyms:	MORG1
Summary:	This gene encodes a member of the WD-40 protein family. The protein is proposed to function as a molecular scaffold for various multimeric protein complexes. The protein associates with several components of the extracellular signal-regulated kinase (ERK) pathway, and promotes ERK activity in response to serum or other signals. The protein also interacts with egl nine homolog 3 (EGLN3, also known as PHD3) and regulates expression of hypoxia-inducible factor 1, and has been purified as part of the spliceosome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2009]

Product images:



Coomassie blue staining of purified WDR83 protein (Cat# [TP301323]). The protein was produced from HEK293T cells transfected with WDR83 cDNA clone (Cat# [RC201323]) using MegaTran 2.0 (Cat# [TT210002]).