

Product datasheet for **TP301336M**

NR2F6 (NM_005234) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human nuclear receptor subfamily 2, group F, member 6 (NR2F6), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201336 representing NM_005234 Red =Cloning site Green =Tags(s)

MAMVTGGWGGPGGDTNGVDKAGGYPRAAEDDSASPPGAASDAEPGDEERPGLQVDCVWCGDKSSGKHYGV
FTCEGCKSFFKRSIRRNLSTCRSNRDCQIDQHHRNQCYCRLKKCFRVGMRKEAVQRGRIPHSPLGAVA
ASSGSPGSALAAVASGGDLFPGQPVSELIAQLLRAEPYPAAAGRFGAGGGAAGAVLGIDNVCELAARLL
FSTVEWARHAPFFPELPVADQVALLRLSWSELFVLNAAQAALPLHTAPLLAAAGLHAAPMAAERAVAFMD
QVRAFQEVDKLGRLQVDSAIEYGLKAIALFTPACGLSDPAHVESLQEKAQVALTEYVRAQYPSQPQRF
GRLLLRLPALRAVPASLISQLFFMRLVGKTIETLIRDMLLSGSTFNWPYGSQ

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	42.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_005225
Locus ID:	2063



[View online »](#)

UniProt ID: [P10588](#), [F1D8R3](#)

RefSeq Size: 1804

Cytogenetics: 19p13.11


RefSeq ORF: 1212

Synonyms: EAR-2; EAR2; ERBAL2

Summary: Transcription factor predominantly involved in transcriptional repression. Binds to promoter/enhancer response elements that contain the imperfect 5'-AGGTCA-3' direct or inverted repeats with various spacings which are also recognized by other nuclear hormone receptors. Involved in modulation of hormonal responses. Represses transcriptional activity of the lutropin-choriogonadotropic hormone receptor/LHCGR gene, the renin/REN gene and the oxytocin-neurophysin/OXT gene. Represses the triiodothyronine-dependent and -independent transcriptional activity of the thyroid hormone receptor gene in a cell type-specific manner. The corepressing function towards thyroid hormone receptor beta/THRB involves at least in part the inhibition of THRB binding to triiodothyronine response elements (TREs) by NR2F6. Inhibits NFATC transcription factor DNA binding and subsequently its transcriptional activity. Acts as transcriptional repressor of IL-17 expression in Th-17 differentiated CD4(+) T cells and may be involved in induction and/or maintenance of peripheral immunological tolerance and autoimmunity. Involved in development of forebrain circadian clock; is required early in the development of the locus coeruleus (LC).[UniProtKB/Swiss-Prot Function]

Protein Families: Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

Product images:



Coomassie blue staining of purified NR2F6 protein (Cat# [TP301336]). The protein was produced from HEK293T cells transfected with NR2F6 cDNA clone (Cat# [RC201336]) using MegaTran 2.0 (Cat# [TT210002]).