

Product datasheet for TP301391

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

EIF4EL3 (EIF4E2) (NM_004846) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human eukaryotic translation initiation factor 4E family member 2

(EIF4E2), 20 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC201391 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MNNKFDALKDDDSGDHDQNEENSTQKDGEKEKTERDKNQSSSKRKAVVPGPAEHPLQYNYTFWYSRRT

PG

RPTSSQSYEQNIKQIGTFASVEQFWRFYSHMVRPGDLTGHSDFHLFKEGIKPMWEDDANKNGGKWIIRL

R

KGLASRCWENLILAMLGEQFMVGEEICGAVVSVRFQEDIISIWNKTASDQATTARIRDTLRRVLNLPPNT

IMEYKTHTDSIKMPGRLGPQRLLFQNLWKPRLNVP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 28.2 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 004837



EIF4EL3 (EIF4E2) (NM_004846) Human Recombinant Protein - TP301391

Locus ID: 9470

 UniProt ID:
 O60573

 RefSeq Size:
 1078

 Cytogenetics:
 2q37.1

 RefSeq ORF:
 735

Synonyms: 4E-LP; 4EHP; EIF4EL3; h4EHP; IF4e

Summary: Recognizes and binds the 7-methylguanosine-containing mRNA cap during an early step in


the initiation (PubMed:9582349, PubMed:17368478, PubMed:25624349). Acts as a repressor of translation initiation (PubMed:22751931). In contrast to EIF4E, it is unable to bind eIF4G (EIF4G1, EIF4G2 or EIF4G3), suggesting that it acts by competing with EIF4E and block

assembly of eIF4F at the cap (By similarity).[UniProtKB/Swiss-Prot Function]

Protein Families: Transcription Factors

Protein Pathways: Insulin signaling pathway, mTOR signaling pathway

Product images:



Coomassie blue staining of purified EIF4E2 protein (Cat# TP301391). The protein was produced from HEK293T cells transfected with EIF4E2 cDNA clone (Cat# [RC201391]) using MegaTran 2.0 (Cat# [TT210002]).