

Product datasheet for TP301407M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NAP1L4 (NM_005969) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human nucleosome assembly protein 1-like 4 (NAP1L4), 100 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC201407 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MADHSFSDGVPSDSVEAAKNASNTEKLTDQVMQNPRVLAALQERLDNVPHTPSSYIETLPKAVKRRINAL KQLQVRCAHIEAKFYEEVHDLERKYAALYQPLFDKRREFITGDVEPTDAESEWHSENEEEEKLAGDMKSK VVVTEKAAATAEEPDPKGIPEFWFTIFRNVDMLSELVQEYDEPILKHLQDIKVKFSDPGQPMSFVLEFHF EPNDYFTNSVLTKTYKMKSEPDKADPFSFEGPEIVDCDGCTIDWKKGKNVTVKTIKKKQKHKGRGTVRTI TKQVPNESFFNFFNPLKASGDGESLDEDSEFTLASDFEIGHFFRERIVPRAVLYFTGEAIEDDDNFEEGE

EGEEELEGDEEGEDEDDAEINPKV

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 42.6 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeg: NP 005960

Locus ID: 4676



ORIGENE

UniProt ID: Q99733

RefSeq Size: 2564

11p15.4 Cytogenetics: 1125 RefSeq ORF:

Synonyms: hNAP2; NAP1L4b; NAP2; NAP2L

Summary: This gene encodes a member of the nucleosome assembly protein (NAP) family which can

interact with both core and linker histones. It can shuttle between the cytoplasm and nucleus,


suggesting a role as a histone chaperone. This gene is one of several located near the

imprinted gene domain of 11p15.5, an important tumor-suppressor gene region. Alterations in this region have been associated with the Beckwith-Wiedemann syndrome, Wilms tumor,

rhabdomyosarcoma, adrenocortical carcinoma, and lung, ovarian, and breast cancer.

[provided by RefSeq, Jul 2008]

Product images:



Coomassie blue staining of purified NAP1L4 protein (Cat# [TP301407]). The protein was produced from HEK293T cells transfected with NAP1L4 cDNA clone (Cat# [RC201407]) using MegaTran 2.0 (Cat# [TT210002]).