

Product datasheet for TP301445

RPLP0 (NM_053275) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human ribosomal protein, large, P0 (RPLP0), transcript variant 2, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201445 protein sequence Red =Cloning site Green =Tags(s)
	MPREDRATWKSNYFLKIIQLDDYPKCFIVGADNVGSKQMQQIRMSLRGKAVLMGKNTMMRKAIRGH LE NNPALEKLLPHIRGNVGFVFTKEDLTEIRDMLLANKVPAARAGAIAPCEVTVPAQNTGLGPEKTSFFQA LGITTKISRGTIEILSDVQLIKTGDKVGASEATLLNMLNISPFSFGLVIQVFDNGSIYNPEVLDITEET LHSRFLEGVRNVASVCLQIGYPTVASVPHSIINGYKRVLALSVDYTFPLAEKVKAFADPSAFVAAAP VAAATTAAPAAAAAPAKVEAKEESESEDEDMGFGLFD
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	34.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_444505




[View online »](#)

Locus ID:	6175
UniProt ID:	P05388
RefSeq Size:	1289
Cytogenetics:	12q24.23
RefSeq ORF:	951
Synonyms:	L10E; LP0; P0; PRLP0; RPP0

Summary: Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein, which is the functional equivalent of the E. coli L10 ribosomal protein, belongs to the L10P family of ribosomal proteins. It is a neutral phosphoprotein with a C-terminal end that is nearly identical to the C-terminal ends of the acidic ribosomal phosphoproteins P1 and P2. The P0 protein can interact with P1 and P2 to form a pentameric complex consisting of P1 and P2 dimers, and a P0 monomer. The protein is located in the cytoplasm. Transcript variants derived from alternative splicing exist; they encode the same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Jul 2008]

Protein Pathways: Ribosome

Product images:



Coomassie blue staining of purified RPLP0 protein (Cat# TP301445). The protein was produced from HEK293T cells transfected with RPLP0 cDNA clone (Cat# [RC201445]) using MegaTran 2.0 (Cat# [TT210002]).