

Product datasheet for TP301493M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HSPC210 (GSKIP) (NM_016472) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human chromosome 14 open reading frame 129 (C14orf129), 100 μg

Species: Human Expression Host: HEK293T

Expression cDNA Clone >RC201493 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

METDCNPMELSSMSGFEEGSELNGFEGTDMKDMRLEAEAVVNDVLFAVNNMFVSKSLRCADDVAYINVET KERNRYCLELTEAGLKVVGYAFDQVDDHLQTPYHETVYSLLDTLSPAYREAFGNALLQRLEALKRDGQS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 15.5 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 057556

 Locus ID:
 51527

 UniProt ID:
 Q9P0R6

 RefSeq Size:
 2251

 Cytogenetics:
 14q32.2






RefSeq ORF: 417

Synonyms: C14orf129; HSPC210

Summary: This gene encodes a protein that is involved as a negative regulator of GSK3-beta in the Wnt

signaling pathway. The encoded protein may play a role in the retinoic acid signaling pathway by regulating the functional interactions between GSK3-beta, beta-catenin and cyclin D1, and it regulates the beta-catenin/N-cadherin pool. The encoded protein contains a GSK3-beta interacting domain (GID) in its C-terminus, which is similar to the GID of Axin. The protein also contains an evolutionarily conserved RII-binding domain, which facilitates binding with protein kinase-A and GSK3-beta, enabling its role as an A-kinase anchoring protein. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, Dec 2012]

Product images:



Coomassie blue staining of purified GSKIP protein (Cat# [TP301493]). The protein was produced from HEK293T cells transfected with GSKIP cDNA clone (Cat# [RC201493]) using MegaTran 2.0 (Cat# [TT210002]).