

Product datasheet for TP301518

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

STAP2 (NM_001013841) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human signal transducing adaptor family member 2 (STAP2),

transcript variant 2, 20 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC201518 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MASALRPPRVPKPKGVLPSHYYESFLEKKGPCDRDYKKFWAGLQGLTIYFYNSNRDFQHVEKLNLGAFEK LTDEIPWGSSRDPGTHFSLILRNQEIKFKVETLECREMWKGFILTVVELRVPTDLTLLPGHLYMMSEVLA KEEARRALETPSCFLKVSRLEAQLLLERYPECGNLLLRPSGDGADGVSVTTRQMHNGTHVVRHYKVKREG PKYVIDVEQPFSCTSLDAVVNYFVSHTKKALVPFLLDEDYEKVLGYVEADKENGENVWVAPSAPGPGPAP CTGGPKPLSPASSQDKLPPLPPLPNQEENYVTPIGDGPAVDYENQDVASSSWPVILKPKKLPKPPAKLPK

PPVGPKPEPKVFNGGLGRKLPVSSAQPLFPTAGLADMTAELQKKLEKRRALEH

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK Predicted MW: 44.7 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001013863





Synonyms:

Locus ID: 55620

UniProt ID: Q9UGK3

RefSeq Size: 1419

Cytogenetics: 19p13.3

BKS


RefSeq ORF: 1209

Summary: This gene encodes the substrate of breast tumor kinase, an Src-type non-receptor tyrosine

> kinase. The encoded protein possesses domains and several tyrosine phosphorylation sites characteristic of adaptor proteins that mediate the interactions linking proteins involved in signal transduction pathways. Alternative splicing results in multiple transcript variants.

[provided by RefSeq, Jul 2008]

Product images:



Coomassie blue staining of purified STAP2 protein (Cat# TP301518). The protein was produced from HEK293T cells transfected with STAP2 cDNA clone (Cat# [RC201518]) using

MegaTran 2.0 (Cat# [TT210002]).