

## Product datasheet for **TP301518L**

### STAP2 (NM\_001013841) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human signal transducing adaptor family member 2 (STAP2), transcript variant 2, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201518 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MASALRPPRPVKPKGVLP SHYYESFLEKKGPCDRDYKKFWAGLQGLTIYFYNSNRDFQHVEKLNILGAFEK  
LTDEIPWGSSRDPGTHFSLILRNQEIKFKVETLECREMWKGFILTVELRVPTDLTLLPGHLYMMSEVLA  
KEEARRALET PSCFLKVSRLAQ LLLERYPECGNLLLRPSGDGADGVSVTTRQMHNGTHVVRHYKVKREG  
PKYVIDVEQPFSC TSLDAVVNYFVSHTKKALVPFLLEDYEVKLVGYVEADKENGENVWAPSAPGPGPAP  
CTGGPKPLSPASSQDKLPPLPPLPNQEENYVTPIGDGPVDYENQDVASSSWPVILKPKKLPKPPAKLPK  
PPVGPKEPKVFENGGLGRKLPVSSAQPLFPPTAGLADMTAELQKKLEKRRALEH

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**


Tag:	C-Myc/DDK
Predicted MW:	44.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_001013863</a>



[View online »](#)

Locus ID:	55620
UniProt ID:	<a href="#">Q9UGK3</a>
RefSeq Size:	1419
Cytogenetics:	19p13.3
RefSeq ORF:	1209
Synonyms:	BKS
Summary:	This gene encodes the substrate of breast tumor kinase, an Src-type non-receptor tyrosine kinase. The encoded protein possesses domains and several tyrosine phosphorylation sites characteristic of adaptor proteins that mediate the interactions linking proteins involved in signal transduction pathways. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2008]

### Product images:



Coomassie blue staining of purified STAP2 protein (Cat# [TP301518]). The protein was produced from HEK293T cells transfected with STAP2 cDNA clone (Cat# [RC201518]) using MegaTran 2.0 (Cat# [TT210002]).