

Product datasheet for TP301580M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Heat Shock Factor 2 Binding Protein (HSF2BP) (NM 007031) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human heat shock transcription factor 2 binding protein (HSF2BP),

100 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

>RC201580 protein sequence Red=Cloning site Green=Tags(s)

MGEAGAAEEACRHMGTKEEFVKVRKKDLERLTTEVMQIRDFLPRILNGEVLESFQKLKIVEKNLERKEQE LEQLKMDCEHFKARLETVQADNIREKKEKLALRQQLNEAKQQLLQQAEYCTEMGAAACTLLWGVSSSEEV VKAILGGDKALKFFSITGQTMESFVKSLDGDVQELDSDESQFVFALAGIVTNVAAIACGREFLVNSSRVL LDTILQLLGDLKPGQCTKLKVLMLMSLYNVSINLKGLKYISESPGFIPLLWWLLSDPDAEVCLHVLRLVQ

SVVLEPEVFSKSASEFRSSLPLQRILAMSKSRNPRLQTAAQELLEDLRTLEHNV

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-Myc/DDK

Predicted MW: 37.5 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 008962

Locus ID: 11077





RefSeq ORF:

Heat Shock Factor 2 Binding Protein (HSF2BP) (NM_007031) Human Recombinant Protein -TP301580M

UniProt ID: 075031, Q6IAT7

RefSeq Size: 1916 Cytogenetics: 21q22.3

Synonyms: MEILB2; POF19

1002


Summary: HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the

> trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2

activation. [provided by RefSeq, Jul 2008]

Protein Families: Transcription Factors

Product images:



Coomassie blue staining of purified HSF2BP protein (Cat# [TP301580]). The protein was produced from HEK293T cells transfected with HSF2BP cDNA clone (Cat# [RC201580]) using

MegaTran 2.0 (Cat# [TT210002]).