

## Product datasheet for **TP301599L**

### ORC2 (NM\_006190) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human origin recognition complex, subunit 2-like (yeast) (ORC2L), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201599 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MSKPELKEDKMLEVHFVGGDDVNLHILDREGGAKLKKERAQLLVNPKKIIKKPEYDLEEDDQEVLDKQNY  
VEIMGRDVQESLKNKSATGGGNKVYSFQNRKHSEKMAKLASELAKTPQKSVSFLKNDPEITINVPQSSK  
GHSASDKVQPKNNDKSEFLSTAPRSLRKRLIVPRSHSDSESEYSASNSDDEGVAQEHEEDTNAVIFSQK  
IQAQNRVVSAPVGKETPSKRMKRDKTSDLVVEEYFEAHSSSKVLTSDRTLQKLKRAKLDQQTLRNLSSKVS  
PSFSAELKQLNQYKELFHKWMLQLHLGFNIVLYGLGSKRDLLERFRTTMLQDSIHVINGFFPGISVKS  
VLNSITEEVLDMGTFRSILDQLDWIVNKFEDSSLELFLLIHNLDQMLRGEKSQQIIGQLSSLHNIYL  
IASIDHLNAPLMWDHAKQSLFNWLWYETTTSPYTEETSYENSLLVKQSGSLPLSSLTHVLRSLTPNARG  
IFRLLIKYQLDNQDNPSYIGLSFQDFYQQCREAFLVNSDLTLRAQLTEFRDHKLIRTKKGTGDGVEYLLIP  
VDNGTLTDFLEKEEEEEA

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**

Tag:	C-Myc/DDK
Predicted MW:	65.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** [NP\\_006181](#)

**Locus ID:** 4999

**UniProt ID:** [Q13416](#), [A0A024R411](#)

**RefSeq Size:** 3140

**Cytogenetics:** 2q33.1

**RefSeq ORF:** 1731


**Synonyms:** ORC2L

**Summary:** The origin recognition complex (ORC) is a highly conserved six subunits protein complex essential for the initiation of the DNA replication in eukaryotic cells. Studies in yeast demonstrated that ORC binds specifically to origins of replication and serves as a platform for the assembly of additional initiation factors such as Cdc6 and Mcm proteins. The protein encoded by this gene is a subunit of the ORC complex. This protein forms a core complex with ORC3, -4, and -5. It also interacts with CDC45 and MCM10, which are proteins known to be important for the initiation of DNA replication. This protein has been demonstrated to specifically associate with the origin of replication of Epstein-Barr virus in human cells, and is thought to be required for DNA replication from viral origin of replication. Alternatively spliced transcript variants have been found, one of which is a nonsense-mediated mRNA decay candidate. [provided by RefSeq, Oct 2010]

**Protein Families:** Stem cell - Pluripotency, Transcription Factors

**Protein Pathways:** Cell cycle

### Product images:



Coomassie blue staining of purified ORC2 protein (Cat# [TP301599]). The protein was produced from HEK293T cells transfected with ORC2 cDNA clone (Cat# [RC201599]) using MegaTran 2.0 (Cat# [TT210002]).