

Product datasheet for TP301652L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RTF2 (NM 016407) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human chromosome 20 open reading frame 43 (C20orf43), 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC201652 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MGCDGGTIPKRHELVKGPKKVEKVDKDAELVAQWNYCTLSQEILRRPIVACELGRLYNKDAVIEFLLDKS AEKALGKAASHIKSIKNVTELKLSDNPAWEGDKGNTKGDKHDDLQRARFICPVVGLEMNGRHRFCFLRCC GCVFSERALKEIKAEVCHTCGAAFQEDDVIVLNGTKEDVDVLKTRMEERRLRAKLEKKTKKPKAAESVSK PDVSEEAPGPSKVKTGKPEEASLDSREKKTNLAPKSTAMNESSSGKAGKPPCGATKRSIADSEESEAYKS

LFTTHSSAKRSKEESAHWVTHTSYCF

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 33.7 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 057491

Locus ID: 51507



RTF2 (NM_016407) Human Recombinant Protein - TP301652L

UniProt ID: Q9BY42
RefSeq Size: 1672

Cytogenetics: 20q13.31

RefSeq ORF: 918


Synonyms: C20orf43; CDAO5; HSPC164; RTFDC1; SHUJUN-3

Summary: Replication termination factor which is a component of the elongating replisome (Probable).

Required for ATR pathway signaling upon DNA damage and has a positive activity during DNA replication. Might function to facilitate fork pausing at replication fork barriers like the rDNA. May be globally required to stimulate ATR signaling after the fork stalls or encounters a lesion (Probable). Interacts with nascent DNA (PubMed:29290612).[UniProtKB/Swiss-Prot

Function]

Product images:



Coomassie blue staining of purified RTF2 protein (Cat# [TP301652]). The protein was produced from HEK293T cells transfected with RTF2 cDNA clone (Cat# [RC201652]) using MegaTran 2.0 (Cat# [TT210002]).