

Product datasheet for TP301652M

RTF2 (NM_016407) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins Recombinant protein of human chromosome 20 open reading frame 43 (C20orf43), 100 µg **Description:** Species: Human HEK293T **Expression Host: Expression cDNA Clone** >RC201652 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s) MGCDGGTIPKRHELVKGPKKVEKVDKDAELVAQWNYCTLSQEILRRPIVACELGRLYNKDAVIEFLLDKS AEKALGKAASHIKSIKNVTELKLSDNPAWEGDKGNTKGDKHDDLQRARFICPVVGLEMNGRHRFCFLRCC GCVFSERALKEIKAEVCHTCGAAFQEDDVIVLNGTKEDVDVLKTRMEERRLRAKLEKKTKKPKAAESVSK PDVSEEAPGPSKVKTGKPEEASLDSREKKTNLAPKSTAMNESSSGKAGKPPCGATKRSIADSEESEAYKS LFTTHSSAKRSKEESAHWVTHTSYCF **TRTRPLEQKLISEEDLAANDILDYKDDDDKV** C-Myc/DDK Tag: Predicted MW: 33.7 kDa **Concentration:** >0.05 µg/µL as determined by microplate BCA method > 80% as determined by SDS-PAGE and Coomassie blue staining **Purity: Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol Recombinant protein was captured through anti-DDK affinity column followed by **Preparation:** conventional chromatography steps. For testing in cell culture applications, please filter before use. Note that you may experience Note: some loss of protein during the filtration process. Store at -80°C. Storage: Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles. RefSeq: NP 057491 Locus ID: 51507



liew online


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	RTF2 (NM_016407) Human Recombinant Protein – TP301652M
UniProt ID:	<u>Q9BY42</u>
RefSeq Size:	1672
Cytogenetics:	20q13.31
RefSeq ORF:	918
Synonyms:	C20orf43; CDAO5; HSPC164; RTFDC1; SHUJUN-3
Summary:	Replication termination factor which is a component of the elongating replisome (Probable). Required for ATR pathway signaling upon DNA damage and has a positive activity during DNA replication. Might function to facilitate fork pausing at replication fork barriers like the rDNA. May be globally required to stimulate ATR signaling after the fork stalls or encounters a lesion (Probable). Interacts with nascent DNA (PubMed:29290612).[UniProtKB/Swiss-Prot Function]

Product images:



Coomassie blue staining of purified RTF2 protein (Cat# [TP301652]). The protein was produced from HEK293T cells transfected with RTF2 cDNA clone (Cat# [RC201652]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US