

Product datasheet for **TP301655L**

RFC3 (NM_002915) Human Recombinant Protein

Product data:

| | |
|--|---|
| Product Type: | Recombinant Proteins |
| Description: | Recombinant protein of human replication factor C (activator 1) 3, 38kDa (RFC3), transcript variant 1, 1 mg |
| Species: | Human |
| Expression Host: | HEK293T |
| Expression cDNA Clone or AA Sequence: | >RC201655 protein sequence Red =Cloning site Green =Tags(s) |

MSLWVDKYRPCSLGRLDYHKEQAAQLRNLVQCGDFPHLLVYGPSGAGKKTRIMCILRELYGVGVEKLRIE
HQTITTPSKKKIEISTIASNYHLEVNPSDAGNSDRVIQEMLKTVAQSQQLETNSQRDFKVVLLTEVDKL
TKDAQHALRRTMEKYMSTCRLILCCNSTSKVIPPIRSRCLAVRVPAPSIEDICHVLSTVCKKEGLNLPSQ
LAHRLAEKSCRNLRKALLMCEACRVQQYPFTADQEIPETDWEVYLRETANAIVSQQTPQRLLLEVRGRLYE
LLTHCIPPEIIMKGLLSELLHNCDGQLKGEVAQMAAYYEHRLQLGSKAIYHLEAFVAKFMALYKFKMEDG
LEGMMF

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

| | |
|-----------------------|--|
| Tag: | C-Myc/DDK |
| Predicted MW: | 40.4 kDa |
| Concentration: | >0.05 µg/µL as determined by microplate BCA method |
| Purity: | > 80% as determined by SDS-PAGE and Coomassie blue staining |
| Buffer: | 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol |
| Preparation: | Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps. |
| Note: | For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. |
| Storage: | Store at -80°C. |
| Stability: | Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles. |
| RefSeq: | <u>NP_002906</u> |



[View online »](#)


| | |
|---------------|---|
| Locus ID: | 5983 |
| UniProt ID: | P40938 , A0A024RDQ8 |
| RefSeq Size: | 2396 |
| Cytogenetics: | 13q13.2 |
| RefSeq ORF: | 1068 |
| Synonyms: | RFC38 |

Summary: The elongation of primed DNA templates by DNA polymerase delta and DNA polymerase epsilon requires the accessory proteins proliferating cell nuclear antigen (PCNA) and replication factor C (RFC). RFC, also named activator 1, is a protein complex consisting of five distinct subunits of 140, 40, 38, 37, and 36 kDa. This gene encodes the 38 kDa subunit. This subunit is essential for the interaction between the 140 kDa subunit and the core complex that consists of the 36, 37, and 40 kDa subunits. Alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008]

Protein Families: Stem cell - Pluripotency

Protein Pathways: DNA replication, Mismatch repair, Nucleotide excision repair

Product images:



Coomassie blue staining of purified RFC3 protein (Cat# [TP301655]). The protein was produced from HEK293T cells transfected with RFC3 cDNA clone (Cat# [RC201655]) using MegaTran 2.0 (Cat# [TT210002]).