

Product datasheet for TP301725

HYPE (FICD) (NM_007076) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human FIC domain containing (FICD), 20 µg

Species: Human

Expression Host: HEK293T

Expression cDNA Clone or AA Sequence: >RC201725 protein sequence
Red=Cloning site Green=Tags(s)

MMLIPMASVMAVTEPKWVSVSRFLWVTLSSMVLGSLALLLPLGAVEEQCLAVLKGLYLLRSKPDRQHAATKCTSPSTELSITSRGATLLVAKTKASPAGKLEARAALNQALEMKRQKREKAQKLFMHALKMDPDFVDALTEFGIFSEEDKDIIQADYLYTRALTISPYHEKALVNRDRTLPLVEEIDQRYFSIIDSKVKKVMSIPKGNSALRRVMEETYHHIYHTVAIEGNTLTLSEIRHILETRYAVPGKSLEEQNEVIGMHAAMKYINTTLVSRIGSVTISDVLEIHRRVLGYVDPVEAGRFRTTQVLVGHHPHPQDVEKQMQEFVQWLNSEEAMNLHPVEFAALAHYKLVYIHPFIDGNGRTSRLLMNLILMQAGYPPITIRKEQRSYYHVLEAANEGDVRPFIRFIAKCTETTLDTLLFATTEYSVALPEAQPNSHSGFKETLPVKP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 51.6 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_009007](#)




[View online »](#)

Locus ID: 11153
UniProt ID: [Q9BVA6](#), [A0A024RBM8](#)
RefSeq Size: 1651
Cytogenetics: 12q23.3
RefSeq ORF: 1374
Synonyms: HIP13; HYPE; UNQ3041

Summary: Protein that can both mediate the addition of adenosine 5'-monophosphate (AMP) to specific residues of target proteins (AMPylation), and the removal of the same modification from target proteins (de-AMPylation), depending on the context (By similarity). The side chain of Glu-231 determines which of the two opposing activities (AMPyase or de-AMPyase) will take place (By similarity). Acts as a key regulator of the ERN1/IRE1-mediated unfolded protein response (UPR) by mediating AMPylation or de-AMPylation of HSPA5/BiP (PubMed:25601083). In unstressed cells, acts as an adenylyltransferase by mediating AMPylation of HSPA5/BiP at 'Thr-518', thereby inactivating it (By similarity). In response to endoplasmic reticulum stress, acts as a phosphodiesterase by mediating removal of ATP (de-AMPylation) from HSPA5/BiP at 'Thr-518', leading to restore HSPA5/BiP activity (By similarity). Although it is able to AMPylate RhoA, Rac and Cdc42 Rho GTPases in vitro, Rho GTPases do not constitute physiological substrates (PubMed:19362538, PubMed:25601083).[UniProtKB/Swiss-Prot Function]

Protein Families: Transmembrane

Product images:



Coomassie blue staining of purified FICD protein (Cat# TP301725). The protein was produced from HEK293T cells transfected with FICD cDNA clone (Cat# [RC201725]) using MegaTran 2.0 (Cat# [TT210002]).