

## Product datasheet for TP301745

### HDAC1 (NM\_004964) Human Recombinant Protein

#### Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human histone deacetylase 1 (HDAC1), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201745 representing NM_004964 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	<p>MAQTQGTTRRKVCYYYDGDVGNYYYGQGHMPKPHRIRMTHNLLLNYGLYRKMEIYRPHKANAEEMTKYHSD DYIKFLRSIRPDNMSEYSKQMQRFNVGEDCPVFDGLFEFCQLSTGGSVASAVKLNKQQTDIAVNWAGGLH HAKKSEASGFCYVNDIVLAILELLKYHQRVLYIDIDIHHDGVEEAFYTTDRVMTVSFHKYGEYFPGTGD LRDIGAGKGKYYAVNYPLRDGIDDESIEAIFKPVMSKVMEMFQPSAVVLQCGSDSLSGDRLGCFNLTIKG HAKCVEFVKSFNLPMLMLGGGGYTIRNVARCWYETAVALDTEIPNELPYNDYFEYFGPDFKLHISPSNM TNQNTNEYLEKIKQRLFENLRMLPHAPGVQMQAIPEDAIEESGDEDEDDEDPDKRISICSSDKRIACEEEF SDSEEEGEGGRKNSSNFKKAKRVKTEDEKEKDPEEKKEVTEEEKTKEEKPEAKGVKEEVKLA</p> <p><b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b></p>
Tag:	C-Myc/DDK
Predicted MW:	54.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_004955</u>



View online »

<b>Locus ID:</b>	3065
<b>UniProt ID:</b>	<a href="#">Q13547</a> , <a href="#">Q6IT96</a>
<b>RefSeq Size:</b>	2091
<b>Cytogenetics:</b>	1p35.2-p35.1
<b>RefSeq ORF:</b>	1446
<b>Synonyms:</b>	GON-10; HD1; KDAC1; RPD3; RPD3L1
<b>Summary:</b>	Histone acetylation and deacetylation, catalyzed by multisubunit complexes, play a key role in the regulation of eukaryotic gene expression. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex. It also interacts with retinoblastoma tumor-suppressor protein and this complex is a key element in the control of cell proliferation and differentiation. Together with metastasis-associated protein-2, it deacetylates p53 and modulates its effect on cell growth and apoptosis. [provided by RefSeq, Jul 2008]
<b>Protein Families:</b>	Adult stem cells, Druggable Genome, Stem cell - Pluripotency, Stem cell relevant signaling - DSL/Notch pathway, Transcription Factors
<b>Protein Pathways:</b>	Cell cycle, Chronic myeloid leukemia, Huntington's disease, Notch signaling pathway, Pathways in cancer

### Product images:



Coomassie blue staining of purified HDAC1 protein (Cat# TP301745). The protein was produced from HEK293T cells transfected with HDAC1 cDNA clone (Cat# [RC201745]) using MegaTran 2.0 (Cat# [TT210002]).