

Product datasheet for **TP301755**

CBS (NM_000071) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human cystathionine-beta-synthase (CBS), 20 µg

Species: Human

Expression Host: HEK293T

**Expression cDNA Clone
or AA Sequence:** >RC201755 protein sequence
Red=Cloning site **Green**=Tags(s)

MPSETPQAEVGP TGCPHRSGPHSAKGSLEKGSPEDEKEAKEPLWIRPDAPSRCTWQLGRPASESPHHHTPP
AKSPKILPDILKKIGDTPMVRINKIGKKFGLKCELLAKCEFFNAGGSVKDRISLRMIEDAERDGTLPKPGD
TIIPTSGNTGIGLALAAAVRGYRCIIVMPEKMSSEKVDVLRALGAEIVRTPNARFSDPESHVGVAVRL
KNEIPNSHILDQYRNASNPLAHYDTTAEILQQC DGKLDMLVASVGTGGTITGIARKLKEKCPGCRIGV
DPEGSILAEPEELNQTEQTTYEVEGIGYDFIPTVLDRTVVDKWFKSNDEEAFTFARMLIAQEGLLCGGSA
GSTVAVAVKAAQELQEGQRCVVILPDSVRNYMTKFLSDRWMLQKGFLEEDLTEKKPWWWHLRVQELGLS
APLTVLPTITCGHTIILREKGFQAPVVDEAGVILGMVTLGNMLSSLLAGKVQPSDQVGVKVIYKQFKQI
RLTDTLGRLSHILEMDHFALVWHEQIQYHSTGKSSQRQMVFGVTAIDLLNFVAAQERDQK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 60.4 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online >](#)

RefSeq: [NP_000062](#)

Locus ID: 875

UniProt ID: [P35520](#), [P0DN79](#), [Q9NTF0](#)

RefSeq Size: 2609

Cytogenetics: 21q22.3

RefSeq ORF: 1653


Synonyms: CBSL; HIP4

Summary: The protein encoded by this gene acts as a homotetramer to catalyze the conversion of homocysteine to cystathionine, the first step in the transsulfuration pathway. The encoded protein is allosterically activated by adenosyl-methionine and uses pyridoxal phosphate as a cofactor. Defects in this gene can cause cystathionine beta-synthase deficiency (CBS), which can lead to homocystinuria. This gene is a major contributor to cellular hydrogen sulfide production. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Feb 2016]

Protein Families: Druggable Genome

Protein Pathways: Cysteine and methionine metabolism, Glycine, serine and threonine metabolism, Metabolic pathways, Selenoamino acid metabolism

Product images:



Coomassie blue staining of purified CBS protein (Cat# TP301755). The protein was produced from HEK293T cells transfected with CBS cDNA clone (Cat# [RC201755]) using MegaTran 2.0 (Cat# [TT210002]).