

Product datasheet for **TP301755L**

CBS (NM_000071) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human cystathionine-beta-synthase (CBS), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201755 protein sequence Red=Cloning site Green=Tags(s)

MPSETPQAEVGP TGCPHRSGPHSAKGSLEKGS PEDKEAKEPLWIRPDAPSRCTWQLGRPASESPHHHTPP
AKSPKILPDILKKIGDTPMVRINKIGKKFGLKCELLAKCEFFNAGGSVKDRISLRMIEDAERDGTLPKPGD
TIIPTSGNTGIGLALAAAVRGYRCIIVMPEKMSSEKVDVLRALGAEIVRTPTNARFSDPESHVGVAVRL
KNEIPNSHILDQYRNASNPLAHYDTTAEILQQC DGKLDMLVASVGTGGTITGIARKLKEKCPGCR IIGV
DPEGSILAEPEELNQTEQTTYEVEGIGYDFIPTVLDR TVVDKWFKSNDEEAFTFARMLIAQEGLLCGGSA
GSTVAVAVKAAQELQEGQRCVVILPDSVRNYMTKFLSDRWMLQKGF LKEEDLTEKKPWWWHLRVQELGLS
APLTVLPTITCGHTIILREKGFDPVDEAGVILGMVTLGNMLSSLLAGKVQPSDQVGVKVIYKQFKQI
RLTDTLGR LSHILEMDHFALVWHEQIQYHSTGKSSQRQMVFGVVT AIDLLNFVAAQERDQK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	60.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP_000062](#)

Locus ID: 875

UniProt ID: [P35520](#), [P0DN79](#), [Q9NTF0](#)

RefSeq Size: 2609

Cytogenetics: 21q22.3

RefSeq ORF: 1653


Synonyms: CBSL; HIP4

Summary: The protein encoded by this gene acts as a homotetramer to catalyze the conversion of homocysteine to cystathionine, the first step in the transsulfuration pathway. The encoded protein is allosterically activated by adenosyl-methionine and uses pyridoxal phosphate as a cofactor. Defects in this gene can cause cystathionine beta-synthase deficiency (CBS), which can lead to homocystinuria. This gene is a major contributor to cellular hydrogen sulfide production. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Feb 2016]

Protein Families: Druggable Genome

Protein Pathways: Cysteine and methionine metabolism, Glycine, serine and threonine metabolism, Metabolic pathways, Selenoamino acid metabolism

Product images:



Coomassie blue staining of purified CBS protein (Cat# [TP301755]). The protein was produced from HEK293T cells transfected with CBS cDNA clone (Cat# [RC201755]) using MegaTran 2.0 (Cat# [TT210002]).