

## Product datasheet for **TP301828**

### SMAD4 (NM\_005359) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human SMAD family member 4 (SMAD4), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201828 representing NM_005359 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MDNMSITNTPTSNDACLSIVHSLMCHRQGGSETFAKRAIESLVKKLKEKKDELDSLITAITTNGAHP  
CVTIQRTLDGRLQVAGRKGFPHVIYARLWRWPDHLKKNELKHVKYCQYAFDLKCDSCVNPYHYERVSP  
G  
IDLSGLTLQSNAPSSMMVKDEYVHDFEGQPSLSTEGHSIQTIQHPPSNRASTETYSTPALLAPSES  
NATS TANFPNIPVASTSQPASILGGSHSEGLLQIASGPQPGQQQNGFTGQPATYHHNSTTTWTGSRT  
APYTPNL PHHQNGHLQHPPMPPHPGHYWPVHNELAFQPPISNHPAPEYWCSIAFYFEMDVQVGET  
FKVPSSCPIV TV  
DGYVDPSSGGDRFCLGQLSNVHRTEAIERARLHIGKGVQLECKGEGDVWVRLSDHAVFVQSY  
LDREAGR APGDAVHKIYPSAYIKVFDLRQCHRQMQQAATAQAAAAAQA AA VAGNIPGPGSVGGI  
APAISLSAAAGI GVDDLRLCILRMSFVKGWGPDYPRQSIKETPCWIEIHLHRLQLLDEVLHTMPI  
ADPQPLD

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**


Tag:	C-Myc/DDK
Predicted MW:	60.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Bioactivity:	Co-immunoprecipitation (PMID: <a href="#">26772959</a> )
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.



[View online »](#)

<b>Storage:</b>	Store at -80°C.
<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_005350</a>
<b>Locus ID:</b>	4089
<b>UniProt ID:</b>	<a href="#">Q13485</a>
<b>RefSeq Size:</b>	3220
<b>Cytogenetics:</b>	18q21.2
<b>RefSeq ORF:</b>	1656
<b>Synonyms:</b>	DPC4; JIP; MADH4; MYHRS
<b>Summary:</b>	<p>This gene encodes a member of the Smad family of signal transduction proteins. Smad proteins are phosphorylated and activated by transmembrane serine-threonine receptor kinases in response to transforming growth factor (TGF)-beta signaling. The product of this gene forms homomeric complexes and heteromeric complexes with other activated Smad proteins, which then accumulate in the nucleus and regulate the transcription of target genes. This protein binds to DNA and recognizes an 8-bp palindromic sequence (GTCTAGAC) called the Smad-binding element (SBE). The protein acts as a tumor suppressor and inhibits epithelial cell proliferation. It may also have an inhibitory effect on tumors by reducing angiogenesis and increasing blood vessel hyperpermeability. The encoded protein is a crucial component of the bone morphogenetic protein signaling pathway. The Smad proteins are subject to complex regulation by post-translational modifications. Mutations or deletions in this gene have been shown to result in pancreatic cancer, juvenile polyposis syndrome, and hereditary hemorrhagic telangiectasia syndrome. [provided by RefSeq, Aug 2017]</p>
<b>Protein Families:</b>	Druggable Genome, Transcription Factors
<b>Protein Pathways:</b>	Adherens junction, Cell cycle, Chronic myeloid leukemia, Colorectal cancer, Pancreatic cancer, Pathways in cancer, TGF-beta signaling pathway, Wnt signaling pathway

### Product images:



Coomassie blue staining of purified SMAD4 protein (Cat# TP301828). The protein was produced from HEK293T cells transfected with SMAD4 cDNA clone (Cat# [RC201828]) using MegaTran 2.0 (Cat# [TT210002]).