

Product datasheet for **TP301872L**

ASCC1 (NM_015947) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human activating signal cointegrator 1 complex subunit 1 (ASCC1), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC201872 protein sequence Red =Cloning site Green =Tags(s)
	 MEVLRPQLIRIDGRNYRKNPQVEQTYQHEEDEEDFYQGSMECADEPCDAYEVEQTPQGFRSTLRAPSLLY KHIVGKRGDTRKKIEMETKTSISIPKPGQDGEIVITGQHRNGVISARTRIDVLLDTFRRKQPFTHFLAFF LNEVEVQEGFLRFQEEVLAKCSMDHGVDSIFQNPKKLHLTIGMLVLLSEEEIQQTCEMLQQCKEEFIND ISGGKPLEVEMAGIEYMNDPFGMVDVLYAKVHMKDGSNRLQELVDRVLERFQASGLIVKEWNSVKLHATV MNTLFRKDPNAEGRYNLYTAEGKYIFKERESFDGRNLIKLFENFYFGSLKLNLSIHISQRFTVDSFGNYAS CGQIDFS TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	41 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_057031</u>



[View online »](#)

Locus ID:	51008
UniProt ID:	Q8N9N2
RefSeq Size:	2150
Cytogenetics:	10q22.1
RefSeq ORF:	1071
Synonyms:	ASC1p50; CGI-18; RP11-150D20.4

Summary: This gene encodes a subunit of the activating signal cointegrator 1 (ASC-1) complex. The ASC-1 complex is a transcriptional coactivator that plays an important role in gene transactivation by multiple transcription factors including activating protein 1 (AP-1), nuclear factor kappa-B (NF-kB) and serum response factor (SRF). The encoded protein contains an N-terminal KH-type RNA-binding motif which is required for AP-1 transactivation by the ASC-1 complex. Mutations in this gene are associated with Barrett esophagus and esophageal adenocarcinoma. Alternatively spliced transcripts encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2011]

Product images:



Coomassie blue staining of purified ASC1 protein (Cat# [TP301872]). The protein was produced from HEK293T cells transfected with ASC1 cDNA clone (Cat# [RC201872]) using MegaTran 2.0 (Cat# [TT210002]).