

Product datasheet for TP302032M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TBCA (NM_004607) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human tubulin folding cofactor A (TBCA), 100 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC202032 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MADPRVRQIKIKTGVVKRLVKEKVMYEKEAKQQEEKIEKMRAEDGENYDIKKQAEILQESRMMIPDCQRR

LEAAYLDLQRILENEKDLEEAEEYKEARLVLDSVKLEA

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-Myc/DDK
Predicted MW: 12.7 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 004598

Locus ID: 6902

UniProt ID: <u>075347</u>

RefSeq Size: 679

Cytogenetics: 5q14.1






RefSeq ORF:

324

Summary:

The product of this gene is one of four proteins (cofactors A, D, E, and C) involved in the pathway leading to correctly folded beta-tubulin from folding intermediates. Cofactors A and D are believed to play a role in capturing and stabilizing beta-tubulin intermediates in a quasi-native confirmation. Cofactor E binds to the cofactor D/beta-tubulin complex; interaction with cofactor C then causes the release of beta-tubulin polypeptides that are committed to the native state. This gene encodes chaperonin cofactor A. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2014]

Product images:



Coomassie blue staining of purified TBCA protein (Cat# [TP302032]). The protein was produced from HEK293T cells transfected with TBCA cDNA clone (Cat# [RC202032]) using MegaTran 2.0 (Cat# [TT210002]).