

Product datasheet for TP302059M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Caspase-7 (CASP7) (NM 033339) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human caspase 7, apoptosis-related cysteine peptidase (CASP7),

transcript variant gamma, 100 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone > or AA Sequence:

>RC202059 protein sequence Red=Cloning site Green=Tags(s)

MADEQGCIEEQGVEDSANEDSVDAKPDRSSFVPSLFSKKKKNVTMRSIKTTRDRVPTYQYNMNFEKLGKC IIINNKNFDKVTGMGVRNGTDKDAEALFKCFRSLGFDVIVYNDCSCAKMQDLLKKASEEDHTNAACFACI LLSHGEENVIYGKDGVTPIKDLTAHFRGDRCKTLLEKPKLFFIQACRGTELDDGIQADSGPINDTDANPR YKIPVEADFLFAYSTVPGYYSWRSPGRGSWFVQALCSILEEHGKDLEIMQILTRVNDRVARHFESQSDDP

HFHEKKQIPCVVSMLTKELYFSQ

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 34.1 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 203125

Locus ID: 840



ORÏGENE

UniProt ID: P55210

RefSeq Size: 2681

10q25.3 Cytogenetics:

909 RefSeq ORF:

Synonyms: CASP-7; CMH-1; ICE-LAP3; LICE2; MCH3


Summary: This gene encodes a member of the cysteine-aspartic acid protease (caspase) family.

> Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. The precursor of the encoded protein is cleaved by caspase 3 and 10, is activated upon cell death stimuli and induces apoptosis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, May

2012]

Protein Families: Druggable Genome, Protease **Protein Pathways:** Alzheimer's disease, Apoptosis

Product images:



Coomassie blue staining of purified CASP7 protein (Cat# [TP302059]). The protein was produced from HEK293T cells transfected with CASP7 cDNA clone (Cat# [RC202059]) using MegaTran 2.0 (Cat# [TT210002]).