

Product datasheet for TP302105M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

BPGM (NM_199186) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human 2,3-bisphosphoglycerate mutase (BPGM), transcript variant 2,

100 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC202105 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MSKYKLIMLRHGEGAWNKENRFCSWVDQKLNSEGMEEARNCGKQLKALNFEFDLVFTSVLNRSIHTAWLI LEELGQEWVPVESSWRLNERHYGALIGLNREQMALNHGEEQVRLWRRSYNVTPPPIEESHPYYQEIYNDR RYKVCDVPLDQLPRSESLKDVLERLLPYWNERIAPEVLRGKTILISAHGNSSRALLKHLEGISDEDIINI

TLPTGVPILLELDENLRAVGPHQFLGDQEAIQAAIKKVEDQGKVKQAKK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 29.8 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 954655

Locus ID: 669





UniProt ID: <u>P07738</u>, <u>A0A024R782</u>

RefSeq Size: 2121 Cytogenetics: 7q33 RefSeq ORF: 777

Synonyms: DPGM; ECYT8


Summary: 2,3-diphosphoglycerate (2,3-DPG) is a small molecule found at high concentrations in red

blood cells where it binds to and decreases the oxygen affinity of hemoglobin. This gene encodes a multifunctional enzyme that catalyzes 2,3-DPG synthesis via its synthetase activity, and 2,3-DPG degradation via its phosphatase activity. The enzyme also has phosphoglycerate phosphomutase activity. Deficiency of this enzyme increases the affinity of cells for oxygen. Mutations in this gene result in hemolytic anemia. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Sep 2009]

Protein Families: Druggable Genome

Protein Pathways: Glycolysis / Gluconeogenesis, Metabolic pathways

Product images:



Coomassie blue staining of purified BPGM protein (Cat# [TP302105]). The protein was produced from HEK293T cells transfected with BPGM cDNA clone (Cat# [RC202105]) using MegaTran 2.0 (Cat# [TT210002]).