

Product datasheet for TP302127M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CPSF7 (NM 024811) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human pre-mRNA cleavage factor I, 59 kDa subunit (FLJ12529),

transcript variant 1, 100 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone or AA Sequence:

>RC202127 protein sequence Red=Cloning site Green=Tags(s)

MSEGVDLIDIYADEEFNQDPEFNNTDQIDLYDDVLTATSQPSDDRSSSTEPPPPVRQEPSPKPNNKTPAI LYTYSGLRNRRAAVYVGSFSWWTTDQQLIQVIRSIGVYDVVELKFAENRANGQSKGYAEVVVASENSVHK LLELLPGKVLNGEKVDVRPATRQNLSQFEAQARKRECVRVPRGGIPPRAHSRDSSDSADGRATPSENLVP SSARVDKPPSVLPYFNRPPSALPLMGLPPPPIPPPPPLSSSFGVPPPPPGIHYQHLMPPPPRLPPHLAVP PPGAIPPALHLNPAFFPPPNATVGPPPDTYMKASAPYNHHGSRDSGPPPSTVSEAEFEDIMKRNRAISSS AISKAVSGASAGDYSDAIETLLTAIAVIKQSRVANDERCRVLISSLKDCLHGIEAKSYSVGASGSSSRKR

HRSRERSPSRSRESSRRHRDLLHNEDRHDDYFQERNREHERHRDRERDRHH

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-Myc/DDK
Predicted MW: 51.9 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.





RefSeq: NP 079087

 Locus ID:
 79869

 UniProt ID:
 Q8N684

 RefSeq Size:
 3764


Cytogenetics: 11q12.2
RefSeq ORF: 1413
Synonyms: CFIm59

Summary: Cleavage factor Im (CFIm) is one of six factors necessary for correct cleavage and

polyadenylation of pre-mRNAs. CFIm is composed of three different subunits of 25, 59, and 68 kDa, and it functions as a heterotetramer, with a dimer of the 25 kDa subunit binding to two of the 59 or 68 kDa subunits. The protein encoded by this gene represents the 59 kDa subunit, which can interact with the splicing factor U2 snRNP Auxiliary Factor (U2AF) 65 to link

the splicing and polyadenylation complexes. [provided by RefSeq, Oct 2016]

Product images:



Coomassie blue staining of purified CPSF7 protein (Cat# [TP302127]). The protein was produced from HEK293T cells transfected with CPSF7 cDNA clone (Cat# [RC202127]) using MegaTran 2.0 (Cat# [TT210002]).