

Product datasheet for TP302195L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cystathionase (CTH) (NM_001902) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human cystathionase (cystathionine gamma-lyase) (CTH), transcript

variant 1, 1 mg

Species: Human Expression Host: HEK293T

Expression cDNA Clone >RC202195 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MQEKDASSQGFLPHFQHFATQAIHVGQDPEQWTSRAVVPPISLSTTFKQGAPGQHSGFEYSRSGNPTRNC LEKAVAALDGAKYCLAFASGLAATVTITHLLKAGDQIICMDDVYGGTNRYFRQVASEFGLKISFVDCSKI KLLEAAITPETKLVWIETPTNPTQKVIDIEGCAHIVHKHGDIILVVDNTFMSPYFQRPLALGADISMYSA TKYMNGHSDVVMGLVSVNCESLHNRLRFLQNSLGAVPSPIDCYLCNRGLKTLHVRMEKHFKNGMAVAQFL ESNPWVEKVIYPGLPSHPQHELVKRQCTGCTGMVTFYIKGTLQHAEIFLKNLKLFTLAESLGGFESLAEL

PAIMTHASVLKNDRDVLGISDTLIRLSVGLEDEEDLLEDLDQALKAAHPPSGSHS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 44.3 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001893



Cystathionase (CTH) (NM_001902) Human Recombinant Protein - TP302195L

Locus ID: 1491

UniProt ID:P32929RefSeq Size:2140Cytogenetics:1p31.1RefSeq ORF:1215

Summary: This gene encodes a cytoplasmic enzyme in the trans-sulfuration pathway that converts


cystathione derived from methionine into cysteine. Glutathione synthesis in the liver is dependent upon the availability of cysteine. Mutations in this gene cause cystathioninuria. Alternative splicing of this gene results in three transcript variants encoding different isoforms.

[provided by RefSeq, Jun 2010]

Protein Pathways: Cysteine and methionine metabolism, Glycine, serine and threonine metabolism, Metabolic

pathways, Nitrogen metabolism, Selenoamino acid metabolism

Product images:



Coomassie blue staining of purified CTH protein (Cat# [TP302195]). The protein was produced from HEK293T cells transfected with CTH cDNA clone (Cat# [RC202195]) using MegaTran 2.0 (Cat# [TT210002]).