

## **Product datasheet for TP302253**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MYD88 (NM\_002468) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human myeloid differentiation primary response gene (88) (MYD88),

20 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

>RC202253 protein sequence Red=Cloning site Green=Tags(s)

MAAGGPGAGSAAPVSSTSSLPLAALNMRVRRRLSLFLNVRTQVAADWTALAEEMDFEYLEIRQLETQADP TGRLLDAWQGRPGASVGRLLELLTKLGRDDVLLELGPSIEEDCQKYILKQQQEEAEKPLQVAAVDSSVPR TAELAGITTLDDPLGHMPERFDAFICYCPSDIQFVQEMIRQLEQTNYRLKLCVSDRDVLPGTCVWSIASE LIEKRCRRMVVVVSDDYLQSKECDFQTKFALSLSPGAHQKRLIPIKYKAMKKEFPSILRFITVCDYTNPC

**TKSWFWTRLAKALSLP** 

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK

Predicted MW: 33.1 kDa

Concentration:  $>0.05 \mu g/\mu L$  as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeg: NP 002459

**Locus ID:** 4615





UniProt ID: Q99836

RefSeq Size: 2862 Cytogenetics: 3p22.2 RefSeq ORF: 888

Synonyms: IMD68; MYD88D


**Summary:** This gene encodes a cytosolic adapter protein that plays a central role in the innate and

adaptive immune response. This protein functions as an essential signal transducer in the interleukin-1 and Toll-like receptor signaling pathways. These pathways regulate that activation of numerous proinflammatory genes. The encoded protein consists of an N-terminal death domain and a C-terminal Toll-interleukin1 receptor domain. Patients with defects in this gene have an increased susceptibility to pyogenic bacterial infections. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Feb 2010]

**Protein Families:** Druggable Genome

**Protein Pathways:** Apoptosis, Toll-like receptor signaling pathway

## **Product images:**



Coomassie blue staining of purified MYD88 protein (Cat# TP302253). The protein was produced from HEK293T cells transfected with MYD88 cDNA clone (Cat# [RC202253]) using MegaTran 2.0 (Cat# [TT210002]).