

Product datasheet for TP302513

KEAP1 (NM_012289) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human kelch-like ECH-associated protein 1 (KEAP1), transcript variant 2, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC202513 protein sequence Red =Cloning site Green =Tags(s)
	<p>MQPDPRPSGAGACCRFLPLQSQCEGAGDAVMYASTECKAEVTPSQHGNRTFSYTLLEDHTKQAFGIMNEL RLSQQLCDVTLQVKYQDAPAAQFMAHKVVLASSPVFKAMFTNGLREQGMEVVSIEGIHPKVMERLIEFA YTASISMGEKCVLHVMNGAVMYQIDSVVRACSDFLVQQLDPSNAIGIANFAEQIGCVLHQRAREIYMH FGEVAKQEEFFNLSHCQLVTLISRDDLNVRCSEV FHACINWVKYDCEQRRFYVQALLRAVRCHSLTPNF LQMQLQKCEILQSDSRCKDYLVKIFEELTLHKPTQVMPCRAPKVGRLIYTAGGYFRQSLSYLEAYNPSDG TWLRLADLQVPRSLAGCVVGGLLYAVGGRNNSPDGNTDSSALDCYNPMTNQWSPCAPMSVPRNRIGVGV IDGHIYAVGGSHGCIHHNSVERYEPERDEWHLVAPMLTRRIGVGVAVLNRLLYAVGGFDGTNRLNSAECY YPERNEWRMITAMNTIRSGAGVCVLHNCIYAAGGYDGDQLNSVERYDVETETWTFVAPMKHRRSALGIT VHQGRIYVLGGYDGHFTFLDSVECYDPD TDTWSEVTRMTSGRSGVGVAVTMEPCRKQIDQONCTC</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-Myc/DDK
Predicted MW:	69.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_036421](#)

Locus ID: 9817

UniProt ID: [Q14145](#), [A0A024R7C0](#)

RefSeq Size: 2577

Cytogenetics: 19p13.2

RefSeq ORF: 1872


Synonyms: INrf2; KLHL19

Summary: This gene encodes a protein containing KELCH-1 like domains, as well as a BTB/POZ domain. Kelch-like ECH-associated protein 1 interacts with NF-E2-related factor 2 in a redox-sensitive manner and the dissociation of the proteins in the cytoplasm is followed by transportation of NF-E2-related factor 2 to the nucleus. This interaction results in the expression of the catalytic subunit of gamma-glutamylcysteine synthetase. Two alternatively spliced transcript variants encoding the same isoform have been found for this gene. [provided by RefSeq, Jul 2008]

Protein Families: Transcription Factors

Protein Pathways: Ubiquitin mediated proteolysis

Product images:



Coomassie blue staining of purified KEAP1 protein (Cat# TP302513). The protein was produced from HEK293T cells transfected with KEAP1 cDNA clone (Cat# [RC202513]) using MegaTran 2.0 (Cat# [TT210002]).