

## Product datasheet for **TP302594**

### **SNX15 (NM\_013306) Human Recombinant Protein**

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human sorting nexin 15 (SNX15), transcript variant A, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC202594 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	 MSRQAKDDFLRHVTVSDPRTHPKGYTEYKVTAQFISKKDPEDVKEVWVKRYSDFRKLHGDLAYTHRNLFRLEEFPAFPRAQVFGFRFEASVIEERRKGAEDLLRFTVHIPALNNSPQLKEFFRGGEVTRPLEVSRDLHILPPPLIPTPPPDDPRLSPLLPAERRGLEEVPVDPSPSSPAQEALDLLNCESTEEASGSPARGPLTEAELALFDPFSKEEGAAPSPHVAELATMEVESARLDQEPWEPGGQEEEEEDGEGGPTPAYLSQATELITQALRDEKAGAYAAALQGYRDGVHVLLQGVPSDPLPARQEGVKKKAAEYLKRAEILRLHLSQLPP  <b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b>
Tag:	C-Myc/DDK
Predicted MW:	38.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u><a href="#">NP_037438</a></u>
Locus ID:	29907




[View online »](#)

UniProt ID: [Q9NRS6](#), [E5KQS5](#)  
RefSeq Size: 1957  
Cytogenetics: 11q13.1  
RefSeq ORF: 1026  
Synonyms: HSAF001435

**Summary:** This gene encodes a member of the sorting nexin family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. Overexpression of this gene results in a decrease in the processing of insulin and hepatocyte growth factor receptors to their mature subunits. This decrease is caused by the mislocalization of furin, the endoprotease responsible for cleavage of insulin and hepatocyte growth factor receptors. This protein is involved in endosomal trafficking from the plasma membrane to recycling endosomes or the trans-Golgi network. Alternative splicing results in multiple transcript variants. Read-through transcription also exists between this gene and the upstream ADP-ribosylation factor-like 2 (ARL2) gene. [provided by RefSeq, Dec 2010]

### Product images:



Coomassie blue staining of purified SNX15 protein (Cat# TP302594). The protein was produced from HEK293T cells transfected with SNX15 cDNA clone (Cat# [RC202594]) using MegaTran 2.0 (Cat# [TT210002]).