

Product datasheet for TP302627

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PTEN (NM_000314) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human phosphatase and tensin homolog (PTEN), 20 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

Tag: C-Myc/DDK

Predicted MW: 47 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

Recombinant protein was produced with TrueORF clone, RC202627.

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 000305

 Locus ID:
 5728

 UniProt ID:
 P60484

 RefSeq Size:
 5572

 Cytogenetics:
 10q23.31

RefSeq ORF: 1209

Synonyms: 10q23del; BZS; CWS1; DEC; GLM2; MHAM; MMAC1; PTEN1; PTENbeta; TEP1



Summary:

This gene was identified as a tumor suppressor that is mutated in a large number of cancers at high frequency. The protein encoded by this gene is a phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase. It contains a tensin like domain as well as a catalytic domain similar to that of the dual specificity protein tyrosine phosphatases. Unlike most of the protein tyrosine phosphatases, this protein preferentially dephosphorylates phosphoinositide substrates. It negatively regulates intracellular levels of phosphatidylinositol-3,4,5-trisphosphate in cells and functions as a tumor suppressor by negatively regulating AKT/PKB signaling pathway. The use of a non-canonical (CUG) upstream initiation site produces a longer isoform that initiates translation with a leucine, and is thought to be preferentially associated with the mitochondrial inner membrane. This longer isoform may help regulate energy metabolism in the mitochondria. A pseudogene of this gene is found on chromosome 9. Alternative splicing and the use of multiple translation start codons results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Feb 2015]


Protein Families:

Druggable Genome, Phosphatase

Protein Pathways:

Endometrial cancer, Focal adhesion, Glioma, Inositol phosphate metabolism, Melanoma, p53 signaling pathway, Pathways in cancer, Phosphatidylinositol signaling system, Prostate cancer, Small cell lung cancer, Tight junction

Product images:



Coomassie blue staining of purified PTEN protein (Cat# TP302627). The protein was produced from HEK293T cells transfected with PTEN cDNA clone (Cat# [RC202627]) using MegaTran 2.0 (Cat# [TT210002]).