

Product datasheet for TP302693M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

C1orf41 (HSPB11) (NM_016126) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human heat shock protein family B (small), member 11 (HSPB11), 100

με

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC202693 protein sequence

or AA Sequence: Red=Cloning site Green=Tags(s)

MRKIDLCLSSEGSEVILATSSDEKHPPENIIDGNPETFWTTTGMFPQEFIICFHKHVRIERLVIQSYFVQ TLKIEKSTSKEPVDFEQWIEKDLVHTEGQLQNEEIVAHDGSATYLRFIIVSAFDHFASVHSVSAEGTVVS

NLSS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 16.1 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 057210

Locus ID: 51668

UniProt ID: Q9Y547





RefSeq Size: 601

Cytogenetics: 1p32.3 RefSeq ORF: 432

Synonyms: C1orf41; FAP232; HSPCO34; IFT25; PP25

Summary: Component of the IFT complex B required for sonic hedgehog/SHH signaling. May mediate

transport of SHH components: required for the export of SMO and PTCH1 receptors out of the cilium and the accumulation of GLI2 at the ciliary tip in response to activation of the SHH pathway, suggesting it is involved in the dynamic transport of SHH signaling molecules within the cilium. Not required for ciliary assembly. Its role in intraflagellar transport is mainly seen


in tissues rich in ciliated cells such as kidney and testis. Essential for male fertility,

spermiogenesis and sperm flagella formation. Plays a role in the early development of the

kidney. May be involved in the regulation of ureteric bud initiation (By similarity).

[UniProtKB/Swiss-Prot Function]

Product images:



Coomassie blue staining of purified HSPB11 protein (Cat# [TP302693]). The protein was produced from HEK293T cells transfected with HSPB11 cDNA clone (Cat# [RC202693]) using MegaTran 2.0 (Cat# [TT210002]).