

# Product datasheet for TP302716M

## IL18 (NM\_001562) Human Recombinant Protein

### **Product data:**

#### **Product Type: Recombinant Proteins** Recombinant protein of human interleukin 18 (interferon-gamma-inducing factor) (IL18), 100 **Description:** μg Species: Human **Expression Host:** HEK293T **Expression cDNA Clone** >RC202716 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s) MAAEPVEDNCINFVAMKFIDNTLYFIAEDDENLESDYFGKLESKLSVIRNLNDQVLFIDQGNRPLFEDMT DSDCRDNAPRTIFIISMYKDSQPRGMAVTISVKCEKISTLSCENKIISFKEMNPPDNIKDTKSDIIFFQR SVPGHDNKMQFESSSYEGYFLACEKERDLFKLILKKEDELGDRSIMFTVQNED **TRTRPLEQKLISEEDLAANDILDYKDDDDKV** Tag: C-Myc/DDK Predicted MW: 22.1 kDa Concentration: >0.05 µg/µL as determined by microplate BCA method > 80% as determined by SDS-PAGE and Coomassie blue staining Purity: **Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol Recombinant protein was captured through anti-DDK affinity column followed by **Preparation:** conventional chromatography steps. Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. Store at -80°C. Storage: Stable for 12 months from the date of receipt of the product under proper storage and Stability: handling conditions. Avoid repeated freeze-thaw cycles. **RefSeq:** NP 001553 3606 Locus ID: **UniProt ID:** Q14116




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	IL18 (NM_001562) Human Recombinant Protein – TP302716M
RefSeq Size:	1163
Cytogenetics:	11q23.1
RefSeq ORF:	579
Synonyms:	IGIF; IL-1g; IL-18; IL1F4
Summary:	The protein encoded by this gene is a proinflammatory cytokine of the IL-1 family that is constitutively found as a precursor within the cytoplasm of a variety of cells including macrophages and keratinocytes. The inactive IL-18 precursor is processed to its active form by caspase-1, and is capable of stimulating interferon gamma production, and of regulating both T helper (Th) 1 and Th2 responses. This cytokine has been implicated in the injury of different organs, and in potentially fatal conditions characterized by a cytokine storm. In humans, IL-18 gene is located on chromosome 11. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Aug 2020]
Protein Families	: Druggable Genome, Secreted Protein
Protein Pathway	<i>is:</i> Cytokine-cytokine receptor interaction, Cytosolic DNA-sensing pathway, NOD-like receptor signaling pathway

## **Product images:**



Coomassie blue staining of purified IL18 protein (Cat# [TP302716]). The protein was produced from HEK293T cells transfected with IL18 cDNA clone (Cat# [RC202716]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US