

Product datasheet for TP302771M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Apolipoprotein CII (APOC2) (NM_000483) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human apolipoprotein C-II (APOC2), 100 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC202771 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MGTRLLPALFLVLLVLGFEVQGTQQPQQDEMPSPTLLTQVKESLSSYWESAKTAAQNLYEKTYLPAVDEK

LRDLYSKSTAAMSTYTGIFTDQVLSVLKGEE

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 8.9 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 000474

Locus ID: 344

UniProt ID: P02655, A0A024R0T9

RefSeq Size: 738

Cytogenetics: 19q13.32





RefSeq ORF: 303


Synonyms: APO-CII; APOC-II

Summary: This gene encodes a lipid-binding protein belonging to the apolipoprotein gene family. The

protein is secreted in plasma where it is a component of very low density lipoprotein. This protein activates the enzyme lipoprotein lipase, which hydrolyzes triglycerides and thus provides free fatty acids for cells. Mutations in this gene cause hyperlipoproteinemia type IB, characterized by hypertriglyceridemia, xanthomas, and increased risk of pancreatitis and early atherosclerosis. This gene is present in a cluster with other related apolipoprotein genes on chromosome 19. Naturally occurring read-through transcription exists between this gene and the neighboring upstream apolipoprotein C-IV (APOC4) gene. [provided by RefSeq, Mar 2011]

Protein Families: Druggable Genome, Secreted Protein

Product images:



Coomassie blue staining of purified APOC2 protein (Cat# [TP302771]). The protein was produced from HEK293T cells transfected with APOC2 cDNA clone (Cat# [RC202771]) using MegaTran 2.0 (Cat# [TT210002]).