

Product datasheet for **TP302784L**

IMPA1 (NM_005536) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human inositol(myo)-1(or 4)-monophosphatase 1 (IMPA1), transcript variant 1, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC202784 protein sequence Red =Cloning site Green =Tags(s)

MADPWQECMDYAVTLARQAGEVCEAIKNEMNVMKSSPVDLVTATDQKVEKMLISSIKEKYP SHSFIGE
ESVAAGEKSILTDNPTWIIDPIDGTTNFVHRFPFVAVSIGFAVNKKIEFGVVYSCVEGKMYTARKGKGAF
CNGQKLQVSQQEDITKSLLVTELGSSRTPETVRMVLSNMEKLCIPVHGIRSVGTAAVNMLCVATGGADA
YYEMGIHCWDVAGAGIIVTEAGGVLM DVTGGPFDLMSRRVIAANNRILAERIAKEIQVIPLQRDDED

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	30 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_005527</u>
Locus ID:	3612



[View online »](#)

UniProt ID: [P29218](#), [A0A024R830](#)

RefSeq Size: 3396

Cytogenetics: 8q21.13

RefSeq ORF: 831


Synonyms: IMP; IMPA; MRT59

Summary: This gene encodes an enzyme that dephosphorylates myo-inositol monophosphate to generate free myo-inositol, a precursor of phosphatidylinositol, and is therefore an important modulator of intracellular signal transduction via the production of the second messengers myoinositol 1,4,5-trisphosphate and diacylglycerol. This enzyme can also use myo-inositol-1,3-diphosphate, myo-inositol-1,4-diphosphate, scyllo-inositol-phosphate, glucose-1-phosphate, glucose-6-phosphate, fructose-1-phosphate, beta-glycerophosphate, and 2'-AMP as substrates. This enzyme shows magnesium-dependent phosphatase activity and is inhibited by therapeutic concentrations of lithium. Inhibition of inositol monophosphate hydrolysis and subsequent depletion of inositol for phosphatidylinositol synthesis may explain the anti-manic and anti-depressive effects of lithium administered to treat bipolar disorder. Alternative splicing results in multiple transcript variants encoding distinct isoforms. A pseudogene of this gene is also present on chromosome 8q21.13. [provided by RefSeq, Dec 2014]

Protein Families: Druggable Genome

Protein Pathways: Inositol phosphate metabolism, Metabolic pathways, Phosphatidylinositol signaling system

Product images:



Coomassie blue staining of purified IMPA1 protein (Cat# [TP302784]). The protein was produced from HEK293T cells transfected with IMPA1 cDNA clone (Cat# [RC202784]) using MegaTran 2.0 (Cat# [TT210002]).