

Product datasheet for **TP302843M**

HIF1AN (NM_017902) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human hypoxia inducible factor 1, alpha subunit inhibitor (HIF1AN), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA	>RC202843 protein sequence
Clone or AA Sequence:	Red=Cloning site Green=Tags(s)

MAATAAEAVASGSGEPREEAGALGPAWDESQLRSYSFPTRPIPRLSQSDPRAEELIENEPEWLTDTNLV
YPALKWDLEYLQENIGNGDFSVYSASTHKFLYDEKKMANFQNFKPRSNREEMKFHEFVEKLQDIQQRGG
EERLYLQQTLDNTVGRKIVMDFLGFNWNWINKQQGKRGWGQLTSNLLLIGMEGNVTPAHYDEQQNFFAQI
KGYKRCILFPPDQFECLYPYPVHHPCDRQSQVDFDNPDYERFPNFQNVVGYETVWGP GDVLYIPMYWWHH
IESLLNGGITITVNFWYKGAPTPKRIEYPLKAHQKVAIMRNIEKMLGEALGNPQEVGPLLNTMIKGRYN

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	40.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_060372
Locus ID:	55662



[View online »](#)

UniProt ID: [Q9NWT6](#)

RefSeq Size: 6928

Cytogenetics: 10q24.31


RefSeq ORF: 1047

Synonyms: FIH1

Summary: Hydroxylates HIF-1 alpha at 'Asn-803' in the C-terminal transactivation domain (CAD). Functions as an oxygen sensor and, under normoxic conditions, the hydroxylation prevents interaction of HIF-1 with transcriptional coactivators including Cbp/p300-interacting transactivator. Involved in transcriptional repression through interaction with HIF1A, VHL and histone deacetylases. Hydroxylates specific Asn residues within ankyrin repeat domains (ARD) of NFKB1, NFKBIA, NOTCH1, ASB4, PPP1R12A and several other ARD-containing proteins. Also hydroxylates Asp and His residues within ARDs of ANK1 and TNKS2, respectively. Negatively regulates NOTCH1 activity, accelerating myogenic differentiation. Positively regulates ASB4 activity, promoting vascular differentiation.[UniProtKB/Swiss-Prot Function]

Protein Families: Druggable Genome

Product images:



Coomassie blue staining of purified HIF1AN protein (Cat# [TP302843]). The protein was produced from HEK293T cells transfected with HIF1AN cDNA clone (Cat# [RC202843]) using MegaTran 2.0 (Cat# [TT210002]).