

Product datasheet for **TP302860L**

BAG3 (NM_004281) Human Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Homo sapiens BCL2-associated athanogene 3 (BAG3), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC202860 representing NM_004281 Red =Cloning site Green =Tags(s)
	<p>MSAATHSPMMQVASGNGDRDPLPPGWEIKIDPQTGWPFVVDHNSRTTTTWNDPRVPSEGPKETPSSANGPS REGSRLPPAREGHPVYPQLRPGYIPIVLHEGAENRQVHPFHVYPQPGMQRFRTAAAAAPQRSQSPLRG MPETTQPKQCGQVAAAAAQQPASHGPERSQSPAASDCSSSSSSASLPSSGRSSLGSHQLPRGYISIPV IHEQNVTRPAAQPSFHQAQKTHYPAQQGEYQTHQPVYHKIQGDDWEPRPLRAASPFSSVQGASSREGSP ARSSTPLHSPSPIRVHTVDRPQQPMTHRETAPVSQPENKPKPGVGPVPELPPGHIPIQVIRKEVDSKP VSQKPPPPSEKVEVKVPPAPVPCPPSPGPSAVPSSPKSVATEERAAPSTAPAEATPPKPGAEAPPKHP GVLKVEAILEKVQGLEQAVDNFEGKKTDDKYLMIIEYLTKELLALDSVDPEGRADVRQARRDGVKRVQTI LEKLEQKAIDVPGVQVYELQPSNLEADQLQAIMEMGAVAADKGGKKNAGNAEDPHTTETQQPEATAAATS NPSSMTDTPGNPAAP</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-Myc/DDK
Predicted MW:	61.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_004272
Locus ID:	9531
UniProt ID:	O95817
RefSeq Size:	2608
Cytogenetics:	10q26.11
RefSeq ORF:	1725
Synonyms:	BAG-3; BIS; CAIR-1; MFM6
Summary:	BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The protein encoded by this gene contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner. [provided by RefSeq, Jul 2008]
Protein Families:	Druggable Genome

Product images:



Coomassie blue staining of purified BAG3 protein (Cat# [TP302860]). The protein was produced from HEK293T cells transfected with BAG3 cDNA clone (Cat# [RC202860]) using MegaTran 2.0 (Cat# [TT210002]).