

## Product datasheet for TP302929

### Vitronectin (VTN) (NM\_000638) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human vitronectin (VTN), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC202929 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MAPLRPLLILALLAWVALADQESCKGRCTEGFNVDKCCQCDLCSYYQSCCTDYTAECKPQVTRGDVFTM  
PEDEYTVYDDGEEKNNATVHEQVGGPSLSDLQAQSKGNPEQTPVLKPEEEAPAPEVGASKPEGIDSRPE  
TLHPGRPQPPAEEELCSGKPFDAFTDLKNGSLFAFRGQYCYELDEKAVRPGYPKLRDVGIEGPIDAAF  
TRINCQGKTYLFKGSQYWRFDGVLDPDYPRNISDGFDPDNDVAALALPAHSYSGRERVYFFKQKQYW  
EYQFQHQPSQEECEGSSLSAVFEHFAMMQRDSWEDIFELLFWGRTSAGTRQPQFISRDWHGVPQVDA  
AM  
AGRIYISGMAPRPSLAKKQFRHRNRKGYRSQRGHSRGRNQNSRRPSRAMWLSLFSSEESNLGANNYDD  
Y  
RMDWLVPATCEPIQSVFFSGDKYYRVNLRTRRVDTVDPPYPRISIAQYWLGCAPAGHL

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**


Tag:	C-Myc/DDK
Predicted MW:	52.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_000629</a>
<b>Locus ID:</b>	7448
<b>UniProt ID:</b>	<a href="#">P04004</a>
<b>RefSeq Size:</b>	1678
<b>Cytogenetics:</b>	17q11.2
<b>RefSeq ORF:</b>	1434
<b>Synonyms:</b>	V75; VN; VNT
<b>Summary:</b>	The protein encoded by this gene functions in part as an adhesive glycoprotein. Differential expression of this protein can promote either cell adhesion or migration as it links cells to the extracellular matrix through a variety of ligands. These ligands include integrins, plasminogen activator inhibitor-1, and urokinase plasminogen activator receptor. This secreted protein can be present in the plasma as a monomer or dimer and forms a multimer in the extracellular matrix of several tissues. This protein also inhibits the membrane-damaging effect of the terminal cytolytic complement pathway and binds to several serpin serine protease inhibitors. This protein can also promote extracellular matrix degradation and thus plays a role in tumorigenesis. It is involved in a variety of other biological processes such as the regulation of the coagulation pathway, wound healing, and tissue remodeling. The heparin-binding domain of this protein give it anti-microbial properties. It is also a lipid binding protein that forms a principal component of high density lipoprotein. [provided by RefSeq, Aug 2020]
<b>Protein Families:</b>	Druggable Genome, Secreted Protein
<b>Protein Pathways:</b>	ECM-receptor interaction, Focal adhesion

### Product images:



Coomassie blue staining of purified VTN protein (Cat# TP302929). The protein was produced from HEK293T cells transfected with VTN cDNA clone (Cat# [RC202929]) using MegaTran 2.0 (Cat# [TT210002]).