

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TP303045M

### CCDC53 (WASHC3) (NM\_016053) Human Recombinant Protein

### **Product data:**


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human coiled-coil domain containing 53 (CCDC53), 100 $\mu g$
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC203045 protein sequence Red=Cloning site Green=Tags(s)
	MDEDGLPLMGSGIDLTKVPAIQQKRTVAFLNQFVVHTVQFLNRFSTVCEEKLADLSLRIQQIETTLNILD AKLSSIPGLDDVTVEVSPLNVTSVTNGAHPEATSEQPQQSSTQDSGLQESEVSAENILTVAKDPRYARYL KMVQVGVPVMAIRNKMISEGLDPDLLERPDAPVPDGESEKTVEESSDSESSFSD
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	21 kDa
Concentration:	>0.05 $\mu$ g/ $\mu$ L as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP 057137</u>
Locus ID:	51019
UniProt ID:	<u>Q9Y3C0</u>
RefSeq Size:	1015



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	CCDC53 (WASHC3) (NM_016053) Human Recombinant Protein – TP303045M
Cytogenetics:	12q23.2
RefSeq ORF:	582
Synonyms:	CCDC53; CGI-116
Summary:	Acts at least in part as component of the WASH core complex whose assembly at the surface of endosomes seems to inhibit WASH nucleation-promoting factor (NPF) activity in recruiting and activating the Arp2/3 complex to induce actin polymerization, and which is involved in regulation of the fission of tubules that serve as transport intermediates during endosome sorting (PubMed:19922875, PubMed:20498093).[UniProtKB/Swiss-Prot Function]

## **Product images:**



Coomassie blue staining of purified WASHC3 protein (Cat# [TP303045]). The protein was produced from HEK293T cells transfected with WASHC3 cDNA clone (Cat# [RC203045]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US