

Product datasheet for TP303052L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DCPS (NM_014026) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human decapping enzyme, scavenger (DCPS), 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC203052 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MADAAPQLGKRKRELDVEEAHAASTEEKEAGVGNGTCAPVRLPFSGFRLQKVLRESARDKIIFLHGKVNE ASEDGDGEDAVVILEKTPFQVEQVAQLLTGSPELQLQFSNDIYSTYHLFPPRQLNDVKTTVVYPATEKHL QKYLRQDLRLIRETGDDYRNITLPHLESQSLSIQWVYNILDKKAEADRIVFENPDPSDGFVLIPDLKWNQ QQLDDLYLIAICHRRGIRSLRDLTPEHLPLLRNILHQGQEAILQRYRMKGDHLRVYLHYLPSYYHLHVHF TALGFEAPGSGVERAHLLAEVIENLECDPRHYQQRTLTFALRADDPLLKLLQEAQQS

C-Myc/DDK

Predicted MW: 38.4 kDa

Tag:

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 054745

Locus ID: 28960



DCPS (NM_014026) Human Recombinant Protein - TP303052L

UniProt ID: <u>Q96C86</u>, <u>A0A384MTI8</u>

RefSeq Size: 1508
Cytogenetics: 11q24.2
RefSeq ORF: 1011

Synonyms: ARS; DCS1; HINT-5; HINT5; HSL1; HSPC015


Summary: This gene encodes a member of the histidine triad family of pyrophosphatases that removes

short mRNA fragments containing the 5′ mRNA cap structure, which appear in the 3′ → 5′ mRNA decay pathway, following deadenylation and exosome-mediated turnover. This enzyme hydrolyzes the triphosphate linkage of the cap structure (7-methylguanosine nucleoside triphosphate) to yield 7-methylguanosine monophosphate and nucleoside diphosphate. It protects the cell from the potentially toxic accumulation of these short, capped mRNA fragments, and regulates the activity of other cap-binding proteins, which are inhibited by their accumulation. It also acts as a transcript-specific modulator of

pre-mRNA splicing and microRNA turnover. [provided by RefSeq, Apr 2017]

Protein Pathways: RNA degradation

Product images:



Coomassie blue staining of purified DCPS protein (Cat# [TP303052]). The protein was produced from HEK293T cells transfected with DCPS cDNA clone (Cat# [RC203052]) using MegaTran 2.0 (Cat# [TT210002]).