

Product datasheet for TP303115

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cystatin A (CSTA) (NM_005213) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human cystatin A (stefin A) (CSTA), 20 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC203115 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MIPGGLSEAKPATPEIQEIVDKVKPQLEEKTNETYGKLEAVQYKTQVVAGTNYYIKVRAGDNKYMHLKVF

KSLPGQNEDLVLTGYQVDKNKDDELTGF

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 10.8 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: <u>NP 005204</u>

 Locus ID:
 1475

 UniProt ID:
 P01040

 RefSeq Size:
 838

 Cytogenetics:
 3q21.1





RefSeq ORF: 294


Synonyms: AREI; PSS4; STF1; STFA

Summary: The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences.

Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins, and kininogens. This gene encodes a stefin that functions as a cysteine protease inhibitor, forming tight complexes with papain and the cathepsins B, H, and L. The protein is one of the precursor proteins of cornified cell envelope in keratinocytes and plays a role in epidermal development and maintenance. Stefins have been proposed as prognostic and diagnostic tools for cancer.

[provided by RefSeq, Jul 2008]

Product images:



Coomassie blue staining of purified CSTA protein (Cat# TP303115). The protein was produced from HEK293T cells transfected with CSTA cDNA clone (Cat# [RC203115]) using MegaTran 2.0 (Cat# [TT210002]).