

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TP303115L

### Cystatin A (CSTA) (NM\_005213) Human Recombinant Protein

### **Product data:**

Product Type:	Recombinant Proteins	
Description:	Recombinant protein of human cystatin A (stefin A) (CSTA), 1 mg	
Species:	Human	
Expression Host:	HEK293T	
Expression cDNA Clone or AA Sequence:	>RC203115 protein sequence Red=Cloning site Green=Tags(s)	
	MIPGGLSEAKPATPEIQEIVDKVKPQLEEKTNETYGKLEAVQYKTQVVAGTNYYIKVRAGDNKYMHLKVF KSLPGQNEDLVLTGYQVDKNKDDELTGF	
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV	
Tag:	C-Myc/DDK	
Predicted MW:	10.8 kDa	
Concentration:	>0.05 µg/µL as determined by microplate BCA method	
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining	
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol	
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.	
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.	
Storage:	Store at -80°C.	
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.	
RefSeq:	<u>NP 005204</u>	
Locus ID:	1475	
UniProt ID:	<u>P01040</u>	
RefSeq Size:	838	
Cytogenetics:	3q21.1	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Cystatin A (CSTA) (NM_005213) Human Recombinant Protein – TP303115L		
RefSeq ORF:	294		
Synonyms:	AREI; PSS4; STF1; STFA		
Summary:	The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins, and kininogens. This gene encodes a stefin that functions as a cysteine protease inhibitor, forming tight complexes with papain and the cathepsins B, H, and L. The protein is one of the precursor proteins of cornified cell envelope in keratinocytes and plays a role in epidermal development and maintenance. Stefins have been proposed as prognostic and diagnostic tools for cancer. [provided by RefSeq, Jul 2008]		

## **Product images:**

188	_	
98	-	
62	_	
49	-	
38	_	
28	_	
17	_	
14	_	-
63	=	

Coomassie blue staining of purified CSTA protein (Cat# [TP303115]). The protein was produced from HEK293T cells transfected with CSTA cDNA clone (Cat# [RC203115]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US