

## Product datasheet for TP303232M

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RBM17 (NM\_032905) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human RNA binding motif protein 17 (RBM17), transcript variant 1,

100 µg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC203232 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MSLYDDLGVETSDSKTEGWSKNFKLLQSQLQVKKAALTQAKSQRTKQSTVLAPVIDLKRGGSSDDRQIVD TPPHVAAGLKDPVPSGFSAGEVLIPLADEYDPMFPNDYEKVVKRQREERQRQRELERQKEIEEREKRRKD RHEASGFARRPDPDSDEDEDYERERRKRSMGGAAIAPPTSLVEKDKELPRDFPYEEDSRPRSQSSKAAIP

PPVYEEQDRPRSPTGPSNSFLANMGGTVAHKIMQKYGFREGQGLGKHEQGLSTALSVEKTSKRGGKIIVG DATEKDASKKSDSNPLTEILKCPTKVVLLRNMVGAGEVDEDLEVETKEECEKYGKVGKCVIFEIPGAPDD

EAVRIFLEFERVESAIKAVVDLNGRYFGGRVVKACFYNLDKFRVLDLAEQV

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK
Predicted MW: 44.8 kDa

The Real

Concentration:  $>0.05 \mu g/\mu L$  as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 116294





**Locus ID:** 84991

UniProt ID: <u>Q96125</u>, <u>Q5W009</u>

RefSeq Size: 3342
Cytogenetics: 10p15.1
RefSeq ORF: 1203
Synonyms: SPF45


**Summary:** This gene encodes an RNA binding protein. The encoded protein is part of the spliceosome

complex and functions in the second catalytic step of mRNA splicing. Alternatively spliced transcript variants have been described. Related pseudogenes exist on chromosomes 9 and

15. [provided by RefSeq, Mar 2009]

**Protein Pathways:** Spliceosome

## **Product images:**



Coomassie blue staining of purified RBM17 protein (Cat# [TP303232]). The protein was produced from HEK293T cells transfected with RBM17 cDNA clone (Cat# [RC203232]) using MegaTran 2.0 (Cat# [TT210002]).