

## Product datasheet for TP303237M

### FMO2 (NM\_001460) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human flavin containing monooxygenase 2 (non-functional) (FMO2), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC203237 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MAKKVAVIGAGVSGLISLKCCVDEGLEPTCFERTEDIGGWRFKENVEDGRASIYQSVTNTSKEMSCFS  
DFPMPEDFPNFLHNSKLLLEYFRIFAKKFDLLKYIQFQTTVLSVRKCPDFSSSGQWKVVTQNGKEQSAVF  
DAVMVCSGHHILPHIPLKSFPGMERFKGQYFHSRQYKHPDGFEGKRILVIGMNGSGDIAVELSKNAAQV  
FISTRHGTWVMSRISEDGYPWDSVFHTRFRSMLRNVLPRTAVKWMIEQQMNRWFNHENYGLEPQNKYIMK  
EPVLNDDVPSRLLCGAIKVKSTVKELTETSAIFEDGTVEENIDVIIFATGYSFSPFLEDVSLVKNENMV  
SLYKYIFPAHLDKSTLACIGLIQPLGSIFPTAELQARWVTRVFKGLCSLPSERTMMMDIIKRNEKRIDL  
GESQSQTLQTNVYDYLDELALALEIGAKPDFCSLLFKDPKLAVERLYFGPCNSY

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**

Tag:	C-Myc/DDK
Predicted MW:	53.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP\\_001451](#)

Locus ID: 2327

UniProt ID: [Q99518](#), [Q5JPC7](#)

RefSeq Size: 5304

Cytogenetics: 1q24.3

RefSeq ORF: 1413

Synonyms: FMO1B1

**Summary:** This gene encodes a flavin-containing monooxygenase family member. It is an NADPH-dependent enzyme that catalyzes the N-oxidation of some primary alkylamines through an N-hydroxylamine intermediate. However, some human populations contain an allele (FMO2\*2A) with a premature stop codon, resulting in a protein that is C-terminally-truncated, has no catalytic activity, and is likely degraded rapidly. This gene is found in a cluster with other related family members on chromosome 1. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2014]

**Protein Pathways:** Drug metabolism - cytochrome P450

### Product images:



Coomassie blue staining of purified FMO2 protein (Cat# [TP303237]). The protein was produced from HEK293T cells transfected with FMO2 cDNA clone (Cat# [RC203237]) using MegaTran 2.0 (Cat# [TT210002]).