

Product datasheet for **TP303276M**

Tropomyosin 3 (TPM3) (NM_152263) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human tropomyosin 3 (TPM3), transcript variant 1, 100 µg

Species: Human

Expression Host: HEK293T

Expression cDNA Clone or AA Sequence: >RC203276 protein sequence
Red=Cloning site **Green**=Tags(s)

MMEAIIKKMQMLKLDKENALDRAEQAEAEQKQAEERSKQLEDELAAMQKCLKGTEDELDKYSEALKDAQE
KLELAEKKAADAEAEVASLNRRIQLVVEELDRAQERLATALQKLEEAEKAADESERGMKVIENRALKDEE
KMELQEIQKAKHIAEEADRKYEEVARKLVIIEGDLERTEERAELAESKCELEELKNVTNNLKSLEA
QAEKYSQKEDKYEEEIKILTDKLEAETRAEFAERSVAKLEKTIDDLEDELYAQKLKYKAISEELDHALN
DMTSI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 32.8 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_689476](#)

Locus ID: 7170




[View online »](#)

UniProt ID: [P06753](#)
RefSeq Size: 7109
Cytogenetics: 1q21.3
RefSeq ORF: 855
Synonyms: CAPM1; CFTD; HEL-189; HEL-S-82p; hscp30; NEM1; OK/SW-cl.5; TM-5; TM3; TM5; TM30; TM30nm; TPM3nu; TPMsk3; TRK

Summary: This gene encodes a member of the tropomyosin family of actin-binding proteins. Tropomyosins are dimers of coiled-coil proteins that provide stability to actin filaments and regulate access of other actin-binding proteins. Mutations in this gene result in autosomal dominant nemaline myopathy and other muscle disorders. This locus is involved in translocations with other loci, including anaplastic lymphoma receptor tyrosine kinase (ALK) and neurotrophic tyrosine kinase receptor type 1 (NTRK1), which result in the formation of fusion proteins that act as oncogenes. There are numerous pseudogenes for this gene on different chromosomes. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2013]

Protein Pathways: Cardiac muscle contraction, Dilated cardiomyopathy, Hypertrophic cardiomyopathy (HCM), Pathways in cancer, Thyroid cancer

Product images:



Coomassie blue staining of purified TPM3 protein (Cat# [TP303276]). The protein was produced from HEK293T cells transfected with TPM3 cDNA clone (Cat# [RC203276]) using MegaTran 2.0 (Cat# [TT210002]).