

Product datasheet for TP303298

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

EIF4A1 (NM 001416) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human eukaryotic translation initiation factor 4A, isoform 1 (EIF4A1),

20 µg

Species: Human **Expression Host:** HEK293T

Expression cDNA Clone

>RC203298 protein sequence

or AA Sequence: Red=Cloning site Green=Tags(s)

> MSASQDSRSRDNGPDGMEPEGVIESNWNEIVDSFDDMNLSESLLRGIYAYGFEKPSAIQQRAILPCIKGY DVIAQAQSGTGKTATFAISILQQIELDLKATQALVLAPTRELAQQIQKVVMALGDYMGASCHACIGGTNV RAEVQKLQMEAPHIIVGTPGRVFDMLNRRYLSPKYIKMFVLDEADEMLSRGFKDQIYDIFQKLNSNTQVV LLSATMPSDVLEVTKKFMRDPIRILVKKEELTLEGIRQFYINVEREEWKLDTLCDLYETLTITQAVIFIN TRRKVDWLTEKMHARDFTVSAMHGDMDQKERDVIMREFRSGSSRVLITTDLLARGIDVQQVSLVINYDLP

TNRENYIHRIGRGGRFGRKGVAINMVTEEDKRTLRDIETFYNTSIEEMPLNVADLI

SGPTRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

C-Myc/DDK Tag:

Predicted MW: 46 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Recombinant protein was captured through anti-DDK affinity column followed by **Preparation:**

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Store at -80°C. Storage:

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001407





Locus ID: 1973

UniProt ID: P60842 RefSeq Size: 1894 Cytogenetics: 17p13.1 RefSeq ORF: 1218

Synonyms: DDX2A; EIF-4A; eIF-4A-I; EIF4A; eIF4A-I

Summary: ATP-dependent RNA helicase which is a subunit of the eIF4F complex involved in cap


recognition and is required for mRNA binding to ribosome. In the current model of

translation initiation, eIF4A unwinds RNA secondary structures in the 5'-UTR of mRNAs which

is necessary to allow efficient binding of the small ribosomal subunit, and subsequent

scanning for the initiator codon.[UniProtKB/Swiss-Prot Function]

Product images:



Coomassie blue staining of purified EIF4A1 protein (Cat# TP303298). The protein was produced from HEK293T cells transfected with EIF4A1 cDNA clone (Cat# [RC203298]) using

MegaTran 2.0 (Cat# [TT210002]).