

## Product datasheet for **TP303298L**

### EIF4A1 (NM\_001416) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human eukaryotic translation initiation factor 4A, isoform 1 (EIF4A1), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC203298 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MSASQDSRSRDNGPDGMEPEGVIESNWNEIVDSFDDMNLSSELLRGIYAYGFEKPSAIQQRAILPCIKGY  
DVIAQAQSGTGKTATFAISILQQIELDLKATQALVLAPTRELAQQIQKVMALGDYMGASCHACIGGTNV  
RAEVQKLQMEAPHIIVGTPGRVFDMLNRRYLSPKYIKMFVLDEADEMLSRGFKDQIYDIFQKLSNTQVW  
LLSATMPSDVLEVTKKFMRDPIRILVKKEELTLEGIRQFYINVEREEWKLDLTLCDLYETLTITQAVIFIN  
TRRKVDWLTEKMHARDFTVSAMHGDMQKRDVIMREFRSGSSRVLITDLLARGIDVQQVSLVINYLDP  
TNRENYIHRIGRGGFRGRKGVAINMVTEEDKRTLRLDIETFYNTSIEEMPLNVADLI

**SGPTRTRPLEQKLISEEDLAANDILDYKDDDDKV**


Tag:	C-Myc/DDK
Predicted MW:	46 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_001407</a>



[View online »](#)

Locus ID:	1973
UniProt ID:	<a href="#">P60842</a>
RefSeq Size:	1894
Cytogenetics:	17p13.1
RefSeq ORF:	1218
Synonyms:	DDX2A; EIF-4A; eIF-4A-I; EIF4A; eIF4A-I
Summary:	ATP-dependent RNA helicase which is a subunit of the eIF4F complex involved in cap recognition and is required for mRNA binding to ribosome. In the current model of translation initiation, eIF4A unwinds RNA secondary structures in the 5'-UTR of mRNAs which is necessary to allow efficient binding of the small ribosomal subunit, and subsequent scanning for the initiator codon.[UniProtKB/Swiss-Prot Function]

### Product images:



Coomassie blue staining of purified EIF4A1 protein (Cat# [TP303298]). The protein was produced from HEK293T cells transfected with EIF4A1 cDNA clone (Cat# [RC203298]) using MegaTran 2.0 (Cat# [TT210002]).