

Product datasheet for **TP303304L**

Cullin 2 (CUL2) (NM_003591) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human cullin 2 (CUL2), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC203304 protein sequence Red=Cloning site Green=Tags(s)

MSLKPRVWDFDETWNKLLTTIKAVVMLEYVERATWDRFSDIYALCVAYPEPLGERLYTETKIFLENHVR
HLHKRVLESEEQVLVVMYHRYWEEYSKGADYMDCLYRYLNTQFIKKNKLTEADLQYGGVDMNEPLMEIG
ELALDMWRKLMVEPLQAILIRMLLREIKNDRGGEDPNQKVIHGVINSFVHVEQYKKKFKPLKFYQEIFESP
FLTETGEYKQEASNLLQESNCSQYMEKVLGRLKDEEIRCRKYLHPSSYTKVIHECQQRMVADHLQFLHA
ECHNIIRQEKKNDMANMYVLLRAVSTGLPHMIQELQNHIDEGLRATSNTQENMPTLFVESVLEVHGKF
VQLINTVLNGDQHFMSALDKALTSVNYREPKSVCAPPELLAKYCDNLLKKSAGKMTENEVEDRLTSFIT
VFKYIDDKDVFQKFYARMLAKRLIHGLSMSMDSEEAMINKLKQACGYEFTSKLHRMYTDM SVSADLNNKF
NNFIKNQD TVIDLGISFQIYVLQAGAWPLTQAPSSTFAIQELESVQMFELFYSQHFSGRKL TWLHYLC
TGEVKMNYLGKPYVAMVTTYQMAVLLAFNSETVSYKELQDSTQMNEKELTKTIKSLLDVKMINHDSEKE
DIDAESSFLNMNFSSKRTKFKITTSMQKDTPQEMEQRSAVDEDRKMYLQAAIVRIMKARKVLRHNALI
QEVISQSRARFNPSISMIKKCIEVLIDKQYIERSQASADEYSYVA

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	86.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_003582
Locus ID:	8453
UniProt ID:	Q13617 , A0A140VKB1
RefSeq Size:	4238
Cytogenetics:	10p11.21
RefSeq ORF:	2235
Summary:	Core component of multiple cullin-RING-based ECS (ElonginB/C-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complexes, which mediate the ubiquitination of target proteins. ECS complexes and ARIH1 collaborate in tandem to mediate ubiquitination of target proteins (PubMed:27565346). May serve as a rigid scaffold in the complex and may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1. The functional specificity of the ECS complex depends on the substrate recognition component. ECS(VHL) mediates the ubiquitination of hypoxia-inducible factor (HIF).[UniProtKB/Swiss-Prot Function]
Protein Families:	Druggable Genome
Protein Pathways:	Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysis

Product images:



Coomassie blue staining of purified CUL2 protein (Cat# [TP303304]). The protein was produced from HEK293T cells transfected with CUL2 cDNA clone (Cat# [RC203304]) using MegaTran 2.0 (Cat# [TT210002]).