

Product datasheet for TP303438

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

p38 (CRK) (NM_005206) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human v-crk sarcoma virus CT10 oncogene homolog (avian) (CRK),

transcript variant I, 20 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC203438 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MAGNFDSEERSSWYWGRLSRQEAVALLQGQRHGVFLVRDSSTSPGDYVLSVSENSRVSHYIINSSGPRPP VPPSPAQPPPGVSPSRLRIGDQEFDSLPALLEFYKIHYLDTTTLIEPVSRSRQGSGVILRQEEAEYVRAL FDFNGNDEEDLPFKKGDILRIRDKPEEQWWNAEDSEGKRGMIPVPYVEKYRPASASVSALIGGR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 22.7 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 005197

Locus ID: 1398

UniProt ID: P46108





RefSeq Size: 3055

Cytogenetics: 17p13.3 RefSeq ORF: 612

Synonyms: CRKII; p38

Summary: This gene encodes a member of an adapter protein family that binds to several tyrosine-

phosphorylated proteins. The product of this gene has several SH2 and SH3 domains (src-homology domains) and is involved in several signaling pathways, recruiting cytoplasmic proteins in the vicinity of tyrosine kinase through SH2-phosphotyrosine interaction. The N-terminal SH2 domain of this protein functions as a positive regulator of transformation whereas the C-terminal SH3 domain functions as a negative regulator of transformation. Two alternative transcripts encoding different isoforms with distinct biological activity have been

described. [provided by RefSeq, Jul 2008]


Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Chemokine signaling pathway, Chronic myeloid leukemia, ErbB signaling pathway, Fc gamma

R-mediated phagocytosis, Focal adhesion, Insulin signaling pathway, MAPK signaling pathway, Neurotrophin signaling pathway, Pathways in cancer, Regulation of actin cytoskeleton, Renal

cell carcinoma

Product images:



Coomassie blue staining of purified CRK protein (Cat# TP303438). The protein was produced from HEK293T cells transfected with CRK cDNA clone (Cat# [RC203438]) using MegaTran 2.0 (Cat# [TT210002]).