

Product datasheet for TP303468L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CAMKK2 (NM_172226) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human calcium/calmodulin-dependent protein kinase kinase 2, beta

(CAMKK2), transcript variant 7, 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

Recombinant protein was produced with TrueORF clone, RC203468.

Tag: C-Myc/DDK

Predicted MW: 59.4 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 757380

Locus ID: 10645

UniProt ID: Q96RR4, A0A024RBP6

RefSeq Size: 4923

Cytogenetics: 12q24.31

RefSeq ORF: 1623

Synonyms: CAMKK; CAMKKB






Summary:

The product of this gene belongs to the Serine/Threonine protein kinase family, and to the Ca(2+)/calmodulin-dependent protein kinase subfamily. The major isoform of this gene plays a role in the calcium/calmodulin-dependent (CaM) kinase cascade by phosphorylating the downstream kinases CaMK1 and CaMK4. Protein products of this gene also phosphorylate AMP-activated protein kinase (AMPK). This gene has its strongest expression in the brain and influences signalling cascades involved with learning and memory, neuronal differentiation and migration, neurite outgrowth, and synapse formation. Alternative splicing results in multiple transcript variants encoding distinct isoforms. The identified isoforms differ in their ability to undergo autophosphorylation and to phosphorylate downstream kinases. [provided by RefSeq, Jul 2012]

Protein Families: Druggable Genome, Protein Kinase, Transcription Factors

Protein Pathways: Adipocytokine signaling pathway

Product images:



Coomassie blue staining of purified CAMKK2 protein (Cat# [TP303468]). The protein was produced from HEK293T cells transfected with CAMKK2 cDNA clone (Cat# [RC203468]) using MegaTran 2.0 (Cat# [TT210002]).