

Product datasheet for TP303480M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Phosphatidic acid phosphatase type 2B (PLPP3) (NM 003713) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human phosphatidic acid phosphatase type 2B (PPAP2B), transcript

variant 1, 100 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC203480 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MQNYKYDKAIVPESKNGGSPALNNNPRRSGSKRVLLICLDLFCLFMAGLPFLIIETSTIKPYHRGFYCND ESIKYPLKTGETINDAVLCAVGIVIAILAIITGEFYRIYYLKKSRSTIQNPYVAALYKQVGCFLFGCAIS QSFTDIAKVSIGRLRPHFLSVCNPDFSQINCSEGYIQNYRCRGDDSKVQEARKSFFSGHASFSMYTMLYL

VLYLQARFTWRGARLLRPLLQFTLIMMAFYTGLSRVSDHKHHPSDVLAGFAQGALVACCIVFFVSDLFKT

KMTLSLPAPAIRKEILSPVDIIDRNNHHNMM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 34.9 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 003704

Locus ID: 8613





Phosphatidic acid phosphatase type 2B (PLPP3) (NM_003713) Human Recombinant Protein – TP303480M

UniProt ID: <u>014495</u>

RefSeq Size: 3324 Cytogenetics: 1p32.2 RefSeq ORF: 933

Synonyms: Dri42; LPP3; PAP2B; PPAP2B; VCIP

Summary: The protein encoded by this gene is a member of the phosphatidic acid phosphatase (PAP)

family. PAPs convert phosphatidic acid to diacylglycerol, and function in de novo synthesis of glycerolipids as well as in receptor-activated signal transduction mediated by phospholipase D. This protein is a membrane glycoprotein localized at the cell plasma membrane. It has been shown to actively hydrolyze extracellular lysophosphatidic acid and short-chain phosphatidic acid. The expression of this gene is found to be enhanced by epidermal growth


factor in Hela cells. [provided by RefSeq, Mar 2010]

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Ether lipid metabolism, Fc gamma R-mediated phagocytosis, Glycerolipid metabolism,

Glycerophospholipid metabolism, Metabolic pathways, Sphingolipid metabolism

Product images:



Coomassie blue staining of purified PLPP3 protein (Cat# [TP303480]). The protein was produced from HEK293T cells transfected with PLPP3 cDNA clone (Cat# [RC203480]) using

MegaTran 2.0 (Cat# [TT210002]).