

Product datasheet for **TP303488**

RGS10 (NM_001005339) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human regulator of G-protein signaling 10 (RGS10), transcript variant 1, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC203488 protein sequence Red =Cloning site Green =Tags(s)

MFNRAVSRLSRKRPPSDIHDSGSSSSSHQSLKSTAKWAASLENLLEDPEGVKRFREFLKKEFSEENVLF
WLACEDFKKMQDKTQMQEKAKEIYMTFLSSKASSQVNVGQSRLENEKILEEPHPLMFQKLQDQIFNLMKY
DSYSRFLKSDLFLKHKRTEEEEEEDLPDAQTAAKRASRIYNT

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	21 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_001005339
Locus ID:	6001
UniProt ID:	O43665



[View online »](#)


RefSeq Size: 910

Cytogenetics: 10q26.11

RefSeq ORF: 543

Summary: Regulator of G protein signaling (RGS) family members are regulatory molecules that act as GTPase activating proteins (GAPs) for G alpha subunits of heterotrimeric G proteins. RGS proteins are able to deactivate G protein subunits of the Gi alpha, Go alpha and Gq alpha subtypes. They drive G proteins into their inactive GDP-bound forms. Regulator of G protein signaling 10 belongs to this family. All RGS proteins share a conserved 120-amino acid sequence termed the RGS domain. This protein associates specifically with the activated forms of the two related G-protein subunits, G-alpha_{i3} and G-alpha_z but fails to interact with the structurally and functionally distinct G-alpha subunits. Regulator of G protein signaling 10 protein is localized in the nucleus. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Product images:



Coomassie blue staining of purified RGS10 protein (Cat# TP303488). The protein was produced from HEK293T cells transfected with RGS10 cDNA clone (Cat# [RC203488]) using MegaTran 2.0 (Cat# [TT210002]).