

Product datasheet for TP303566L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SPSB4 (NM_080862) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human splA/ryanodine receptor domain and SOCS box containing 4

(SPSB4), 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone or AA Sequence:

>RC203566 protein sequence Red=Cloning site Green=Tags(s)

MGQKLSGSLKSVEVREPALRPAKRELRGAEPGRPARLDQLLDMPAAGLAVQLRHAWNPEDRSLNVFVKD

D

DRLTFHRHPVAQSTDGIRGKVGHARGLHAWQINWPARQRGTHAVVGVATARAPLHSVGYTALVGSDAE

SW

GWDLGRSRLYHDGKNQPGVAYPAFLGPDEAFALPDSLLVVLDMDEGTLSFIVDGQYLGVAFRGLKGKKLY

PVVSAVWGHCEVTMRYINGLDPEPLPLMDLCRRSIRSALGRQRLQDISSLPLPQSLKNYLQYQ

SGPTRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 30 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 543138





Locus ID: 92369

UniProt ID: Q96A44

RefSeq Size: 2455

Cytogenetics: 3q23

RefSeq ORF: 819

Synonyms: SSB-4; SSB4

Summary: Substrate recognition component of a SCF-like ECS (Elongin BC-CUL2/5-SOCS-box protein) E3

ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:21199876, PubMed:15601820). Negatively regulates nitric oxide (NO) production and limits cellular toxicity in activated macrophages by mediating the ubiquitination and proteasomal degradation of NOS2 (PubMed:21199876). Acts as a bridge which links NOS2 with the ECS E3 ubiquitin ligase complex components ELOC and CUL5 (PubMed:21199876). Diminishes EphB2-dependent cell

repulsive responses by mediating the ubiquitination and degradation of EphB2/CTF2


(PubMed:28931592). Regulates cellular clock function by mediating the

ubiquitin/proteasome-dependent degradation of the circadian transcriptional repressor

NR1D1 (PubMed:26392558).[UniProtKB/Swiss-Prot Function]

Protein Families: Druggable Genome

Product images:



Coomassie blue staining of purified SPSB4 protein (Cat# [TP303566]). The protein was produced from HEK293T cells transfected with SPSB4 cDNA clone (Cat# [RC203566]) using MegaTran 2.0 (Cat# [TT210002]).